PLANT DISEASE REPORTER INDEX

Issued by

CROPS RESEARCH DIVISION

Plant Industry Station
Volume 49

Beltsville, Maryland
Number 13

AUTHOR INDEX

ADAMS, ROBERT E., 662
AFANASIEV, M. M., 253
AGRIOS, GEORGE N., 314
AHMED, A. A., (760)
AICHELE, MURIT D., (765), (769), (910)
ALEXANDER, P. M., 258
ALLAM, E. K., 218
ALLAN, R. E., (335)
ALLEN, THOMAS C., Jr., 557
ALMEYDA, N., (550)
ALTMAN, JACK, 600
ANDERSON, GERALD W., 527
ANZALONE, LOUIS, Jr., (757)
ARAGAKI, MINORU, (327), 378
ARMSTRONG, G. M., 412, 673
ARMSTRONG, JOANNE K., (412), (673)
ARNY, D. C., (516)
ASTIN, JOSEPH S., Jr., 68, (360)
AYCOCK, ROBERT, 435

BAGGETT, J. R., (627), 630, (787)
BAJPAI, G. K., (164), (166)
BAKER, A. S., (891)
BANERJEE, A. K., (166)
BARKER, K. R., 47
BARNES, D. K., (676)
BARNES, G. L., 285
BASHAM, J. T., 771
BASU, A. N., (981)
BEAN, GEORGE A., 810
BECKMAN, KENT M., 357
BELL, A. A., 323
BELL, A. H., (849), (982)
BENNETT, C. W., 375
BHATTACHARYA, S. K., (405)
BIRD, G. W., 33, 517
BITTERS, W. P., (7)
BLANK, L. M., 179
BLODGETT, EARLE C., 225, 765, 769, 910
BOHN, G. W., (532)
BOHNENBLUST, K. E., 813
BOKX, J. A. de, 742
BOUDREAUX, H. B., (495)
BOWMAN, DONALD H., (408)
BOYCE, JOHN S., Jr., 322, (698)
BOYLE, LYTTON W., (975)
BRANDSBERG, JOHN, (660)
BRANTLEY, B. B., (879)
BRENNAN, EILEEN, (911)
BRISTER, C. D., (738)
BRITTON, MICHAEL P., 708, (710)
BRODERICK, E. J., (568)
BROWDER, L. E., (308), 455
BROWN, G. ELDON, 490, 646
BRZESKI, M. W., 610
BUMBIERIS, M., (504)
BURLEIGH, JAMES R., (275)
BUTLER, J. D., (708), (710)
BYNUM, H. H., (647)

CADMAN, C. H., 230
CALAVAN, E. C., (668)
CALDWELL, RALPH M., (403)
CALVERT, O. H., 213
CAMPBELL, W. A., 233, (279)
CARPENTER, J. B., 668
CARTER, J. C., (552)
CARVER, R. G., 495
CHANDLER, W. A., 419
CHAPMAN, R. A., (450)
CHARPENTIER, LEON J., (827)
CHI, C. C., 159, 512
CHILDERS, W. R., (512)
CHRISTIANSEN, D. W., (668)
COCHRAN, H. L., (157)
COFFEE, E. G., (603)
COLE, JOHN R., 146, 148, 703
COLLINS, R. J., 865
COMPTON, LEROY E., (403)
CONVERSE, RICHARD, (987)
COON, B. F., (692)
COOPER, W. E., (931)
CORDELL, CHARLES E., 360
COTTER, R. U., (63)
COURTER, J. W., 886
COVEY, RONALD P., Jr., 876
COX, R. S., 1018
CROSSAN, D. F., 312, (748)
CULP, T. W., 433

DAINES, ROBERT H., 300, (911)
DALE, J. L., 202, 757
DAVIS, J. R., (178)
DAVIS, N. D., (931)
DAVIS, SPENCER H., Jr., (835)
DEEMA, NAULCHANTRA H., (478)
DEEP, IRA W., 734
DEMPSEY, A. H., 157
DeWOLFE, T. A., (423)
DICKEY, ROBERT S., 470
DICKSON, R. C., (449)
DIENER, Ul L., 931
DIMALLA, GERALD G., 15
DISHON, I., (154)
DOEPEL, U. L., (943)

DOOLEY, H. L., (581)
DOOLING, O. J., 954
DOWLER, WILLIAM M., (701)
DUKES, P. D., (215), (347), 586, 847
DUNLEAVY, J. M., (339)
DURBIN, R. D., (878), 948

EIDE, CARL J., 638
EISSA, F. M., (598)
EL-HAIDERI, H., (792)
ELLETT, C. WAYNE, 615
ELLIOT, AUTHUR M., (852), (999)
ELSAID, H. M., 18
EL-SHEHEDI, A. A., (760)
ENGLISH, HARLEY, 178
EPPS, JAMES M., 747
EPPS, W. M., (649)
EPSTEIN, ABRAHAM H., (47), 814
ESSER, R. P., 265
EVERETT, PAUL H., (29)

FAIRCHILD, M. L., (213)
FASSULIOTIS, GEORGE, (723)
FATHI, SAAD M., (244)
FELDMAN, A. W., (865)
FENNER, L. M., 360
FENWICK, H. S., (297), (854)
FERRIS, J. M., 69, (72)
FIELDHOUSE, D. J., (748)
FIGARO, PEGGYBETH, (876)
FILER, T. H., Jr., (351), 571, 869
FLEISCHMANN, GEORGE, 132
FORD, RICHARD E., (235), 627, 684, 787
FRAZIER, N. W., 204
FRAZIER, W. A., (630)
FREEMAN, T. E., 382
FREITAG, J. H., (204)
FRIDLUND, PAUL R., 187, 288
FROELICH, R. C., 970
FULTON, N. D., (307)
FUTRELL, MAURICE C., 680

GABRIELSON, R. L., (891)
GALLEGLY, M. E., (233)
GANGULY, B., (981)
GASKIN, T. A., 89, 268, 802, 1017
GAY, J. DAN, 720
GILLESPIE, WILLIAM H., 173
GOH, K. G., (778)
GOLD, A. H., (204)
GOLDEN, A. MORGAN, (5), 987, (991)
GOODING, GUY V., Jr., (797)
GOODMAN, R. N., (380)
GOSS, R. L., (923)
GOTH, R. W., 815, (818)
GOULD, CHARLES J., 149, 443, 923
GRAHAM, T. W., 822
GRAY, L. E., 867
GREELEY, L. W., (575)
GRIFFITHS, D. A., 979
GRUENHAGEN, R. H., 269

GUDAUSKAS, ROBERT T., 91
GUENGERICH, H. W., 39, (380)
GUTHRIE, J. W., 297, 854

HAARD, RICHARD T., (185)
HAGEDORN, D. J., 591
HAGLUND, WILLIAM A., 793
HALEVY, A. H., (775)
HAMILTON, R. I., (253)
HAMPTON, R. O., 235
HAND, LEONARD F., 193
HANSON, E. W., (159)
HARDING, PAUL R., Jr., 332, 965
HARPAZ, I., 34
HARRELL, D. C., (138)
HARRISON, MONTY D., 452
HART, JOHN H., 830, 1016
HAWKSWORTH, FRANK G., (417)
HEALY, MICHAEL J., 710
HEBERT, T. T., (553)
HELTON, A. W., 352
HENDRIX, FLOYD F., Jr., (859)
HENDRIX, J. WALTER, 275
HENSON, LAWRENCE, (450)
HICKEY, K. D., (568)
HILDEBRAND, E. M., 425, 429
HILL, JOHN H., (280), (371)
HILLER, W., (936)
HIMELICK, E. B., (106), 949
HINDI, E., 154
HINDS, THOMAS E., 480, 481
HOFFMANN, J. A., 401
HOLDEMAN, QUINTIN L., (468), 654
HOLTON, C. S., 242, (401)
HOPPE, PAUL E., 516
HORN, N. L., (495)
HORNER, CHESTER E., 393, 538, 581
HOUCK, L. G., 715
HOVELAND, CARL S., (91)
HRUSCHKA, HOWARD W., 959
HUBER, D. M., (297), (854)
HUISINGH, D., 878
HUNTER, J. H., (146)
HUSSIN, HAMID, (734)
HUTCHINS, LEE M., 564

JACKSON, CURTIS R., 928, (931)
JAIN, J. S., (164)
JANSON, BLAIR F., 478, (615)
JARNEVIC, NOEL B., 603
JAWORSKI, C. A., (986)
JEDLINSKI, H., (692)
JENKINS, PHIL, (910)
JENKINS, S. F., Jr., (215), (586), (847)
JENKINS, W. R., (517)
JENSEN, H. J., (509)
JENSEN, RAY E., 975
JERATH, M. L., (729), (875)
JEWELL, F. F., (616)
JHOOTY, J. S., 756
JOFFE, A. Z., 741

JOHN, V. T., 305
JOHNSON, D. E., 898, (900), (903)
JOHNSON, E. M., 450, 590
JOHNSON, HOWARD W., (84), 408, 753
JOHNSON, LOWELL B., 222
JOKELA, J. J., (867)
JONES, JOHN PAUL, 29, 536
JONES, JOHN R., (480)
JUSKA, F. V., (185)

KAHN, ROBERT P., 656
KAIS, A. G., 971
KEASTER, A. J., (213)
KELMAN, ARTHUR, 797, (872)
KENDRICK, E. L., (401), 837, 839, 843
KENNEDY, B. W., (490), (491), (646)
KESSLER, KENNETH J., Jr., 483
KING, T. H., (24)
KINLOCH, B. B., 872
KINMAN, M. L., (383), (433)
KIRKPATRICK, J. D., 7
KLEIN, M., (34)
KLISIEWICZ, J. M., 541
KLOTZ, L. J., 423
KOIRTYOHANN, S. R., (36)
KOLP, B. J., (813)
KONSLER, T. R., (388), (634)
KOZELNICKY, GEORGE M., 207
KRAMER, C. L., (359)
KREITLOW, K. W., 185
KRONSTAD, W. E., (636)
KUHN, C. W., 879

LACKEY, C. F., 1002
LAIRD, E. F., Jr., 449
LAL, S. B., (493)
LAMBE, R. C., 339
LAMEY, H. A., 736, 738
LANGDON, K. R., 12
LATHAM, A. J., 398
LEAR, BERT, (898), 900, 903
LEININGER, L. N., (440)
LEISER, A. T., (69), 72
LEONE, IDA A., 911
LEONG, CHEE KEE, (1011)
LEPPIK, E. E., 940
LIGHTLE, PAUL C., 417
LIM, W. C., (979)
LINCOLN, ARTHUR C., (1007)
LINN, M. B., (398)
LIPSCOMB, HARRIET ANN, 507
LISK, D. J., (568)
LITTRELL, R. H., 78, (536), 649
LIVINGSTON, CLARK H., (452)
LOCKE, SETH BARTON, 546
LORBEER, JAMES W., 522
LOWNSBERY, B. F., 994
LUEPSCHEN, N. S., 392
LUKENS, R. J., 361

MacGOWAN, J. B., (265)
MAHDY, M. T., (170)
MAIER, CHARLES R., 904
MALEKZADEH, F., 439
MARSH, P. B., 138
MARTIN, J. A., (383)
MASSIE, IRA, (590)
MATHUR, R. S., 164, 166
McCLAIN, L. A., 1016
McCRUM, R. C., 958
McDANIEL, M. C., 307
McGRATH, HILDE, 456
McGREW, J. R., 270
McLEAN, D. M., 54
McVEY, DONALD V., 191
MELVIN, CURRIE F., (463)
MEREDITH, DONALD S., 1006
MILBRATH, J. A., (15)
MILHOLLAND, R. D., (435)
MILLER, DOUGLAS R., 647
MILLER, H. N., 51
MILLER, P. M., 1017
MILLER, P. W., 284, 509, 954
MILLER, V. L., (923)
MILLER, WILLIAM E., 268
MILLIKAN, D. F., 36, (39), 380
MINK, G. I., 143
MINTON, NORMAN A., 856
MITCHELL, J. T., (994)
MITTERLING, LLOYD A., 578
MIYAGAWA, S. T., (900), (903)
MOHAMED, HOSNI A., 244, 1013
MOHAMMED, J., (792)
MONROE, R. L., (656)
MORGAN, FREDERIC L., 84, (753)
MORRISON, RICHARD H., 870
MORTON, D. J., (586), (847)
MOSEMAN, J. G., 575
MUELLER, WALTER C., 750
MUKHERJI, S. K., 405
MURPHY, H. C., (692)
MURRAY, EVERETT L., (769)
MUSE, R. R., (323)
MUTHAPPA, B. N., 917

NAGDY, G. A., 170
NATOUR, R. M., 732
NAVARATNAM, S. J., (778), 1011
NEELY, DAN, 106, 552, (949)
NENE, Y. L., (114)
NEVO, D., (154)
NICKLE, W. R., 991
NIELSEN, L. W., 97
NORMAN, C., (529)
NOUR, M. A., (599)

OBRERO, FAUSTINO C., 327
OSHIMA, NAGAYOSHI, (452)
OSTAZESKI, S. A., 855
OTEIFA, B. A., 596, 598
OU, S. H., (127), 778
OUELLETTE, G. B., 909
OWEN, JOHN H., (279), (720), 859

PAI, H. SURENDRANATH, 263
PALTI, J., (741)
PAPASOLOMONTOS, A., 111
PAPAVIZAS, G. C., (256)
PARKER, M. B., 168
PARSONS, J. L., (143)
PARSONS, JACK E., (357)
PASS, HARTWILL, (806)
PATEL, P. N., (591)
PATT, J., 936
PATTERSON, FRED L., (403)
PAVGI, M. S., (58)
PEACHER, P. H., (954)
PEPPER, EVAN H., 519
PERRY, V. G., (51)
PETERSEN, D. H., 701
PETERSON, GLENN W., 121, 124, (269), 529
PETERSON, J. L., 228, 835
PHILLIPPE, MARY RUTH, (74), (210)
PINE, T. S., 109
PODGER, F. D., 943
POKORNY, F. A., (861)
POWELL, W. M., 279, (859), 861
POWELSON, R. L., 636
POWERS, H. R., Jr., 342
PURDY, LAURENCE H., 42, 335

QUEBRAL, F. C., (886)
QUILLIN, RALPH B., (68)
QUINON, VIVENCIO L., (378)

RAABE, ROBERT D., 583, 812, 1015
RAFFRAY, J. B., 500
RAGAB, M. M., 760
RAGHUNATH, T., 916
RAYCHAUDHURI, S. P., 981
REILING, T. P., 24
REINHART, J. H., 60
REXRODE, CHARLES O., 1007
RHOADES, H. L., 259
RIVERA, C. T., 127, (778)
RIVERS, G. W., 383
ROANE, C. W., 665
ROBERTS, B. J., (63)
ROCHOW, W. F., 457, 687, 692
ROMANKO, R. R., 247
RONCADORI, R. W., (972)
ROTHMAN, PAUL G., (408)
RUBIN, H. L., 729, (1016)
RUPPEL, E. G., 550, 676
RUSDEN, P. L., 801
RUSSELL, THOMAS S., (149), (443)

SAVAGE, DONALD C., (332)
SCHAFER, JOHN F., (222), 403
SCHINDLER, A. F., 5
SCHLEHUBER, A. M., 806
SCHMITT, C. G., (367)
SCHNEIDER, C. L., 293
SCHOENEWEISS, DONALD F., 251
SCIARONI, R. H., (900), (903)
SCOTT, H. A., (532)
SENN, L. H., (258)
SHAFIEE, F. A., (598)
SHATLA, M. N., 21, 711
SHEPHERD, ROBERT J., 468, 961
SHER, S. A., 849, 982
SHERROD, L. LYNN, 852, 999
SHIH, MADELINE, (396)
SHOEMAKER, R. A., 832
SHURTLEFF, M. C., (886)
SILBERNAGEL, MATT J., 475, 637
SIMPSON, M. E., (138)
SINCLAIR, J. B., (18), (21), (307), (500), (711)
SINGH, R. P., 493
SINGH, R. S., 114
SINGH, U. P., 58
SITTERLY, W. R., 723
SMILEY, J. H., (590)
SMITH, H. A., (33)
SMITH, WILSON L., Jr., 619, (783)
SMOOT, JOHN J., 463
SNOW, G. A., (970), (971), 972
SOWELL, GROVER, Jr., 605, 607, (879), 895
STALEY, JOHN M., 882
STEGALL, W. A., Jr., (342)
STEWART, DONALD M., 63, 280, 371
STEWART, JAMES L., 456, (481)
STIPES, R. J., (931)
STODDARD, E. M., (1017)
STONE, KENNETH W., (522)
STONER, WARREN N., 918
STRABY, A. E., (832)
STRELECKI, R. F., (228)
STRIDER, D. L., 388, 634
SUMNER, D. R., (529)
SZKOLNIK, MICHAEL, 568

TACHIBANA, H., 396
TAHORI, A. S., 775
TARJAN, A. C., (596)
TAYLOR, JACK, (168)
TAYLOR, L. DAVID, (771)
TAYLOR, W. KENT, 698
TEGETHOFF, ALFRED C., 834
THOMAS, C. A., 119, 256
THOMAS, H. EARL, (1015)
THOMPSON, A. E., (886)
THOMPSON, HUGH E., (359)
THOMPSON, S. S., 215
THORNBERRY, H. H., 74, 210
THYR, B. D., 386
TILL, B. B., (961)
TIMIAN, ROLAND G., 556, 696
TOLER, ROBERT W., 347, 553
TOMPKINS, D. R., 891
TOOLE, E. RICHARD, 81, 351, (869), 986
TROUTMAN, J. L., (665)

TSUE, KING MON, (600)
TU, JUI-CHANG, (275)
TURNER, P. D., 135, 257, 319
TYLER, LEON J., 239

UNNY, K. L., 729, 875
UPADHYAY, JANESHWAR, 724
UPCHURCH, W. J., (36)

VAARTAJA, O., 504
VALLEAU, W. D., (450)
Van PELT, H. M., (265)
VAUGHAN, E. K., (630)

WADDLE, B. M., (138)
WALKER, J. C , 198
WALKER, NELY M., (616)
WALKINSHAW, C. H., 616
WARID, W. A., (760)
WATSON, ALICE, (656)
WATSON, D. R. W., 764
WEATHERS, L. G., 189
WEBB, R. E., 532, 818
WEBSTER, ORRIN J., (680)
WELLS, HOMER D., 487
WESTER, R. E., 447
WHEELER, W. H., (656)
WICKER, ED F., 102
WIEBEL, F. J., 748
WILLIAMS, F. J., 134
WILLIAMS, L. E., (615)
WILLIAMS, PAUL H., (198)
WILLIS, WILLIAM G., 359
WILSON, CHARLES L., 789
WILSON, V. E., 660
WINSTEAD, E. E., (487)
WITCHER, WESLEY, (507)
WOLF, FREDERICK A., 964
WOODBRIDGE, C. G., (891)
WORF, G. L., (47)
WORTHINGTON, JOHN T., III, (619), 783
WUEST, PAUL J., 367
WYCOFF, H. B., (867)
WYSONG, DAVID S., 269

YASSIN, A. M., 599
YOUNG, H. C., Jr., 308, (806)

ZAIGER, D., 565
ZALASKY, H., 50
ZAUMEYER, W. J., (815)
ZEIDLER, G., (775)
ZENTMYER, G. A., (565), (943)
ZIMMER, D. E., (293), 440, 623
ZUMMO, NATALE, 827

SUBJECT INDEX

Abies fraseri: Cylindrocladium scoparium on, 797; new host for Fomes annosus, Polyporus schweinitzii, and Fomes pini, North Carolina, 360
--- lasiocarpa: new host for Scleroderris canker, found in Colorado, 882
Abscission: of petiole bases in stored rhubarb stalks, 959
Acer rubrum: dieback of, 483
--- saccharum: dieback of, 483; occurrence of fungi and bacteria in normal and in discolored heartwood in Ontario, 771
Achlya spp.: 736
Acyrthosiphon porosum: 495
Adenostema fasciculatum: 1002
Agrobacterium rhizogenes: 878
--- tumefaciens: 360; 470; 734
Air pollutants: correlation of weather data with phytotoxicity in New Jersey, 911
Alabama: 91; 832
Alfalfa: Phytophthora root and crown rot of seedlings and mature plants in Mississippi, 753; 21 Rhizoctonia solani isolates from different hosts and locations tested for virulence on, 512; soil culture of, 6 of 8 fertilizer or insecticide amendments used did not harm Xiphinema americanum population levels, 994; viruses of in Rhode Island 750; wilt, pathogenicity studies of 3 forms of Fusarium oxysporum which cause similar symptoms in several varieties, 412
Almond: 596
Alternaria spp.: 138; importance in cotton seedling disease complex in New Mexico, 904
--- cucumerina: 605; 895
--- tenuis: 886
--- tenuissima: cause of new disease of mango in India, 405
Amanita muscaria: 1017
Anthracnose: of bean, 46 domestic and 8 European bean varieties tested for reactions to 4 physiologic races of, 817
---: of cucumber: destructive disease in Florida, promising control, 29; standardization of artificial inoculation methods for obtaining uniform results, 649
---: of guar, caused by Colletotrichum dematium f. truncata, 607
---: of turnip greens, fungicidal control, 419
Antibiotics: cycloheximide sprays fail to

control white pine blister rust at two sites in North Carolina and one in Tennessee, 342; separation of 104 isolates of Agrobacterium tumefaciens into 26 groups on basis of in vitro sensitivity to 14 antibiotics not correlated with the host or virulence, 470
Aphelenchoides ritzema-bosi: 1017
Aphis gossypii: 495; 981; first report as vector of sugarcane mosaic virus, 204
Apple: blister bark, discussion of history, symptoms, distribution, and possible causes of the disease, 765
---: cedar apple rust, 12 clonal rootstock varieties studied for susceptibility to, 578
---: chlorotic leaf spot virus, field spread between adjacent trees suggests direct mechanical leaf to leaf or branch transmission, 958
---: diseases in Kentucky in 1964, 450
---: fruit russet ring virus, first known case of naturally infected McIntosh trees in the United States, 314
---: latent viruses: 'Hopa' crab as indicators for, in Oregon, 15; second season symptoms can result from root grafting of sensitive indicators, 39
---: leaf flecking virus, on McIntosh in Massachusetts, 314
---: powdery mildew, 3 compounds screened at Tree Fruit Experiment Station have mildewcidal properties, 876
---: scab, determination of mercury spray residue on fruit by a sensitive analytical method, neutron activation, and by the Schoniger colorimetric method, 568
---: stem cavity russet, 225
Apricot (see also Prunus): bacterial canker in the southeastern States, 701
---: dieback fungus found on western choke-cherry, 178
Araucaria excelsa: re-evaluation of quarantine significance of Cryptospora longispora on imported seed, 656
Arceuthobium campylopodum: 647
Arizona: 179; 480
Arkansas: 202; 1007
Armillaria mellea: several new hosts for, in California, 812
Articularia quercina var. minor: first report on Quercus dunnii, found in New Mexico, 417
Ascochyta pinodella: 660
Ash: European mountain, previously unreported canker disease was discovered in Illinois in 1963, 251
Aspergillus spp.: 928
--- flavus: invasion of peanut pods and kernels determined at digging time, 931
Atomizer: for inoculating plants with spore-oil suspension, 455
Australia: 504; 943
Azalea: several parasitic nematodes associated with decline in Wisconsin, 47

Bacteria: Agrobacterium tumefaciens, variability in the in vitro sensitivity of 104 isolates to 14 selected antibiotics, 470; crown-gall disease on rhododendron, two records in North America, 360; device for rapid and uniform inoculation of plants with bacteria that require wounds, 878; effect of crop rotation and fertilizers on soil microflora as indicated by fungi and bacteria isolated from roots of cotton seedlings, 1013; gummosis and leaf spotting of sweet cherry appear to be caused by a complex of species conforming to the genera Xanthomonas and Pseudomonas, 380; present in both normal and discolored heartwood of sugar maple in Ontario, 771; possible relationship to hypocotyl collar rot of bean, 134; unidentified seedborne bacterium isolated from 2 watermelon plant introductions which originated in Turkey, 818
Bacterial brown spot: of bean, worst outbreak in Wisconsin in many years during 1964 growing season, 591
--- canker: of stone fruits in the southeastern States, 701; of tomato transplants in the west-central Great Plains, 386; of tomato, description of cotyledonary symptoms, 634
--- crinkle leaf: new disease of soybean produced by unreported bacterial pathogen, 396
--- leaf spot: effect of photoperiod and supplemental nitrogen on susceptibility of sesame, 119
--- spot: of pepper, several chemicals tested in culture and in the field for control, 748
Bacterium tumefaciens: 360
Bark canker: of cacao, first reported epidemic in Caroline Islands, effective control measures, 565
--- measles: of pear, occurrence in Colorado Bartlett orchards in 1962, 392
Barley: leaf rust, identification of 3 new pathogenic strains of Puccinia hordei from 1959-1964, 575
---: powdery mildew, effect of some growth-retarding compounds on, 775
---: yellow dwarf virus, strains grouped into 3 types depending on specificity of aphid transmission, 687
---: stripe mosaic virus, unsuccessful attempt to free seed of virus by heat

treatments, 696
---: stripe rust, possibly first report of stem infection, Oregon, 636
Basal stem rot: of oil palm, results of root inoculations of seedlings in pots with Ganoderma sp., 1011
Bean: alfalfa mosaic virus isolated from snap beans induced enation symptoms in tobacco, 684
---: anthracnose, 46 domestic and 8 European varieties tested for reactions to 4 physiologic races of, 815
---: collar rot of Phaseolus vulgaris 'Tendercrop' possibly related to bacterium, 134
---: curly top virus, differential tolerance of 52 snap varieties to, 475
---: disease survey of bush snap bean producing areas of Washington, Oregon and Idaho during 1964, 637
---: downy mildew, distribution of G strain on lima beans in 1964, 447
---: Fusarium root rot, tests for resistance, 630
---: halo blight: serological detection, 297; and bacterial brown spot in Wisconsin in 1964, worst outbreak in many years, 591
---: Sclerotinia wilt, of green beans, excellent control with Botran when sprayed on lower portion of vines, Washington, 357
---: southern stem rot, effect of some growth-retarding compounds on, 775
---: tobacco ringspot virus, effect of mechanical inoculation of various plant parts on symptom expression, 54
---: yellow mosaic disease, on mung, effect of vector control, 166
Beet (see Sugarbeet)
Bemisia tabaci: 166
Betula alleghaniensis: dieback of, 483
Bioassay: for testing effectiveness of soil chemicals in preventing white rot of onion with an indication of field performance, 546; modified paper disk method used to evaluate 18 fungicides for control of 13 fungi from peanuts, 928; use of cellophane to detect translocation of potential therapeutic chemicals in plant tissue, 949
Birdsfoot trefoil: Phomopsis blight of, host range and testing of Lotus spp. for disease resistance to, 724; first report of natural occurrence of tobacco ringspot virus, 855
Blackline: etiology of disorder in grafted Persian walnuts, tests indicate that no virus is involved, 954
Black pepper (see Pepper)
Black shank: in flue-cured tobacco, spread and control in Georgia since 1959, 215
Black slime: of iris, comparison of broadcast or furrow application with bulb dips in PCNB for control, 443
Blackstem: of cottonwood, studies indicate that it is an infectious disease and that susceptibility may be a heritable trait, 867
Blast disease: of rice, structure of leaves directly related to varietal susceptibility, 170
Blight: of soap-nut trees in India, new disease, 263
---: fire, of pear, testing comparative resistance of Stewart with other Bartlett clones under controlled conditions, 769
---: flower, of camellia, distribution and spread in South Carolina, 258
---: halo: of bean, worst outbreak in Wisconsin in many years during 1964 growing season, 591; (Race 2a), first report on pea , found in Pacific Northwest, 854; serological detection, specific and rapid method for identification of infected plants, 297
---: needle, of Austrian pine, infection and control, 124
---: northern leaf, or corn in Wisconsin in 1964, 516
---: Phomopsis: of birdsfoot trefoil, host range and testing of Lotus spp. for disease resistance, 724; diseased and healthy eastern redcedar seedlings compared for field survival and growth 121; of eastern redcedar seedlings, good control with Puratized Agricultural Spray, 529
---: pod and stem, of soybean, effect on germination and growth in New Jersey, 228
---: seedling, of sugarbeet, 293
---: southern: of annual ryegrass and winter oats, in Mississippi, 408; of iris, 149
---: willow, found in Michigan, first report from central United States, 1016
Blister: new disease of sweetpotato, cause unknown, 97
--- bark: of Red Delicious apple trees, discussion of history, symptoms, distribution, and possible causes of the disease, 765
Blue mold: of tobacco: in Israel, 154; comparison of European and Beltsville isolates of Peronospora tabacina in the greenhouse, 367
Book reviews: "Annual Review of Phytopathology," edited by James G. Horsfall, 456; "Plant Virology," edited by M. K. Corbett and H. D. Sisler, 457; "Strawberry Diseases," by A. G. Plakidas, 270

Botrytis sp.: 619; 783
--- cinerea: 660
Bottlegourd (see Cucurbits)
Broccoli (see Crucifers)
Brussels sprouts (see Crucifers)
Bud union: significance of rootstock selection and tree age in scion overgrowth disorder, 7
Budwood repository: virus-indexed Prunus materials released for research and industry as of August 1964, 188
Bunt: of wheat: control with HCB, PCNB, and TCNA, 42; 70 of 244 spring varieties carry some resistance to key races in screening trials from 1954-62, 839; 38 of 102 winter varieties carry some resistance to key races in screening trials since 1956, 843

Cabbage (see Crucifers)
Cacao: role of polyphenoloxidase as a factor involved in resistance to Phytophthora palmivora, 319
---: bark canker, first reported epidemic in Caroline Islands, effective control measures, 565
---: greenpoint gall, induced by imperfect stage but not by perfect stage of Calonectria rigidiuscula, 564
---: pod rot: behavior of pathogen in soil, 135; ("black pod disease"), effect of spacing and pruning on disease levels, in Ghana, 257
California: 109; 204; 647; 812; 870; 1015
Calonectria rigidiuscula: ability to induce greenpoint gall on cacao during conidial (Fusarium) stage may be lost when perfect (ascospore) stage develops, 564; conidial stage present in necrotic bark from Phytophthora canker infected cacao, 565
Camellia: flower blight, distribution and spread in South Carolina, 258
Cankers: European larch, none found during resurvey of 2826 trees in Massachusetts in 1965, 834; of mountain ash, new disease caused by Fusicoccum sp., 251; on trunks and branches of young Japanese pagoda trees caused by Fusarium lateritium, 835; of tulip poplar caused by Fusarium solani, first report in South Carolina, 507
Cardamon: leaf spot, new disease caused by Sphaceloma sp. in India, 917
Caroline Islands: 565
Carrot-top: stage of Phomopsis disease on western larch, 102
Cassytha filiformis: check list of hosts found in Hawaii in 1963, 583
Castorbean: root and stem rot, susceptibility to Thielaviopsis basicola, 256
Cellophane: used in technique to determine fungistatic activity of translocated compounds in plant tissue, 949
Cephalosporium diospyri: 789
Ceratocystis fagacearum: 68; 269; 954; 1007
--- ulmi: 106; 359; 552; 801
Cercospora spp.: 975
--- plumeriae: found in United States (Florida) for first time, 729
Cercosporella brassicae: 419
Chemicals: amendments used in soil culture of alfalfa did not harm Xiphinema americanum population levels, 994; certain chemical and heat treatments supplement refrigeration for retarding postharvest decay of strawberries, 619: new materials for control of apple powdery mildew, 876; several tested in culture and in the field for control of bacterial spot of pepper, 748
---: biphenyl-resistant mutants of Penicillium digitatum, 965
---: captan reduced decay of stored potato seed pieces, 638
---: chemotherapeutic agents tested for effect on Cucumis virus 2C in bottlegourd, 493
---: cyanogenic compounds produced by a fairy ring fungus, 571
---: DD used as a soil fumigant improved sugarbeet seedling emergence, vigor, and yield, 600
---: dimethyl sulfoxide (DMSO) used with Dithane M-45 and Dyrene for control of Helminthosporium diseases of Kentucky bluegrass, 810
---: Freon-113 as a suspending medium for dispersal of urediospores of Puccinia graminis var. tritici onto wheat, 268
---: growth regulators: tested for effect on 3 fungal diseases and 1 viral disease, 775; gibberellin produced by Fusarium spp. causes "overgrowth" symptoms in Malayan crop plants, 979
---: HCB, PCNB, and TCNA effectively controlled common and dwarf bunts of wheat, 42
---: sodium o-phenylphenate added to a detergent foam and applied to poststorage lemons complemented the action of subsequent biphenyl treatment in packed cartons to reduce decay caused by Penicillium digitatum, 332
---: sulfur dioxide and chemical dips

· tested for control of postharvest decay of red raspberries, 783
---: tetrazolium (TTC) indicates extent of Phytophthora root rot infection of citrus rootstocks in test for resistance or susceptibility, 423
---: urea, effective treatment for stripe smut on 'Merion' bluegrass, 361
Cherry (see also Prunus): dwarfness in Bing caused by virus, may be valuable means of controlling tree size or may present complications, 910; gummosis and leaf spotting symptoms (suggestive of necrotic rusty mottle virus) associated with bacterial infection, 380
---: crown gall, influence of temperature on development in woody hosts, 734
---: latent viruses, prevalent in Washington sweet cherry orchards of 3 age groups, 143
---: "short stem", of 'Lambert' sweet, relationship to other sweet cherry viruses, 253
Choke-cherry: western, apricot dieback fungus found on, 178
Chrysanthemum: wilt of, caused by race 1 of cowpea-wilt Fusarium, not by aster-wilt Fusarium as previously reported, 673
Citrus spp.: 596; hot-water treatment reduced postharvest decay of unethylened oranges as much or more than standard fungicide, 463; pits and protuberances in the xylem of different variety-rootstock combinations in Israel, 936; significance of rootstock selection and tree age in bud union overgrowth disorder of lemon, 7
---: grapefruit virus disease (impietratura) in Cyprus, 111
---: nematodes: apparently not correlated to 4 viruses in Florida groves; and 6 previously unreported species of plant nematodes associated with Florida citrus, 610; burrowing nematode, attempts to improve growth of infected orange trees with undertree drenches of Zinophos, 865; citrus-root nematode, found in citrus groves in Iraq for first time, 1965, 792; plant-parasitic nematodes in established citrus groves in Sudan controlled by applying DBCP in irrigation water, 598
---: Penicillium digitatum: poststorage decay was greatly reduced when a foam washer treatment with sodium o-phenylphenate was applied to fruit before a packed carton biphenyl sheet treatment during shipmant, 332; resistant to biphenyl obtained in culture, 965; warm DCNA reduced decay of lemons during storage, 715
---: Phytophthora root rot, method for estimating degree of resistance or susceptibility of various Rough lemon rootstocks, 423
---: psorosis infection of sour orange rapidly detected by iodine test of leaves, 18
---: stubborn disease, excessive seed abortion in affected fruit, 668
Cladosporium cucumerinum: 29; 312; 388
Climate: factors affecting the spread of hop downy mildew in an arid climate, 247
Clitocybe olearia: associated with decline of oak and sycamore in California, 870
Clover: host-range and pathogenicity studies of Pratylenchus scribneri on white clover and 5 other hosts, 856; 21 Rhizoctonia solani isolates from different sources tested for virulence on red clover, 512; viruses of several varieties in Rhode Island, 750
Colletotrichum dematium f. truncata: 607; 895
--- gloeosporioides: first report of damage to flowering dogwood by, 351
--- gossypii: 138
--- higginsianum: 419
--- lagenarium: 29; 649
--- lindemuthianum: 815
Colorado: 386; 392; 481; 882
Control: of Sclerotinia wilt on greenbeans in Washington with Botran, 357
Coreopsis grandiflora: flower droop associated with unidentified toxicogenic insect, 964
Corn: diseases in Kentucky in 1964, 450
---: maize dwarf mosaic virus: distribution and yield loss in Ohio in 1964, 615; isolations tested for effect on sugarcane, MDMV is possibly a strain of sugarcane mosaic virus, 757; possibly caused by strain of sugarcane mosaic virus, occurrence and transmission in Virginia, 665; screening 93 varieties for susceptibility to, 478
---: Nigrospora cob rot, reaction of 4 open-pollinated varieties to, United Arab Republic, 244
---: northern leaf blight in Wisconsin in 1964, 516
---: stunt virus disease: status in Georgia in 1964, 207; symptoms and distribution in Missouri in 1964, 213
---: sugarcane mosaic virus: recovered

from naturally infected corn, milo maize, and Johnson grass in California, 204; virus disease (Johnson grass strain of sugarcane mosaic) is seedborne in both field and sweet corn, 468
---: virus diseases: in Iowa, mechanically transmitted, 339; in the United States in 1964, 918; distribution, host range, physical characteristics, and aphid transmission studies in Arkansas, 202; new mottle-stunt disease in southern Illinois associated with sap-transmissible virus, 210; tentative key for preliminary field and laboratory decisions, 654
Corrections: 91; 179; 270; 361; 882; 885; 1018
Corynebacterium michiganense: 386; 634
Cotton: effect of chemical seed treatments and varying harvest dates on control of seedling diseases, 720; fungi and bacteria isolated from seedling roots indicate effect of crop rotation and fertilizers on soil microflora, 1013; monitoring duration of critical levels of moisture on plant parts susceptible to disease, 193; reduction in yield caused by diseases in 1964, 307
---: boll rot, observations in South Carolina, 138
---: Fusarium wilt: first occurrence in Arizona, 179; based on isolations, Fusarium wilt was more prevalent than Verticillium wilt in South Plains area of Texas in 1964, 999
---: Rhizoctonia solani: in vitro studies on fungicide tolerance, 500; differences in reaction of 5 varieties to 3 isolates of Rhizoctonia solani suggest existence of physiologic specialization, 711
---: seedling damping-off, in PCNB-treated soil, 21
---: seedling disease complex, 3 Alternaria spp. should be considered important in, New Mexico, 904
---: Verticillium wilt, successful use of disposable, time and space-saving plastic growth pouches which allow plants to be grown under aseptic conditions in studies of penetration by V. albo-atrum into roots, 852
Cottonwood (see Populus)
Cowpea (see Pea)
Crab apple (see Apple)
Crambe abyssinica: virus susceptibility tested for consideration before growing commercially, 74
Cranberry: stylet-bearing nematodes not controlled by bog flooding and low dissolved oxygen concentrations after 2 years, 517
Cronartium coleosporioides (Peridermium harknessii): 527
--- fusiforme: 322; 616; 698; 872; 972; cultures of gall tissue from slash pine were maintained in an apparently viable condition for 5 months, 616
--- quercuum: 527
--- stalactiforme: 909
Crop rotation: effect on soil microflora as indicated by fungi and bacteria isolated from roots of cotton seedlings, United Arab Republic, 1013
Crown gall: influence of temperature on development in woody hosts, 734; on rhododendron, two records in North America, 360
--- rot: Phytophthora, of vetch in Mississippi, 84
--- rust: of oats, racial variability greater in aecial than in uredial collections on buckthorn, 132
Crucifers: anthracnose and white spot, on turnip greens, fungicidal control, 419
---: black rot, evaluation of phenacridane chloride as a disinfectant and seed protectant for control, 439
---: powdery mildew, on cabbage, inheritance of resistance controlled by single dominant gene, 198
---: sugarbeet and cabbage root nematodes: on Brussels sprouts, effectiveness of nematocides in transplant water, 900; effectiveness of soil fumigation, 903
---: sulfur deficiency in field-grown broccoli (first report), description of symptoms in field and greenhouse, 891
Cryptospora longispora: on imported seed of the Norfolk Island pine, re-evaluation of quarantine significance, 656
Cucumber (see Cucurbits)
Cucurbits: effect of mechanical inoculation of various plant parts on symptom expression of TRSV, 54; evaluation of the cucurbita for scab resistance, (mimeographed list of Cucurbita accessions that were screened and are available through Plant Introduction Stations will be supplied if requested), 388; promising results for control of anthracnose, scab, and soil rot in Florida, 29
---: bottlegourd: effect of chemotherapeutic agents on Cucumis virus 2C, 493

---: cucumber: anthracnose, standardization of artificial inoculation methods for obtaining consistently uniform results, 649; scab, fungicide sprays for control, 312
---: squash: isolates of Choanephora cucurbitarum from cowpea pathogenic on, 347; squash mosaic virus disease, study of in Cairo, Egypt, 218
---: watermelon: unidentified seedborne bacterium isolated from 2 plant introductions which originated in Turkey, 818; watermelon mosaic viruses 1 and 2, identified in specimens from production areas of the South and West, 532
Cultures: of fusiform rust-infected slash pine, 616
Curvularia maculans: 928
Cuscuta approximata var. urceolata: 1015
--- sandwichiana: check list of hosts found in Hawaii in 1963, 583
--- subinclusa: 189
Cylindrocladium scoparium: 797
Cyprus: 111
Cytospora canker: on Douglas-fir, recurrence in Colorado, 481
--- chrysosperma: 867
--- kunzei var. kunzei: 481

Daedalea confragosa: 986
Damping-off: of Pinus radiata, promising seed treatments for control in South Australia, 504; pre-emergence, of soybean in Minnesota, 646
Dasyscypha willkommii: 834
Decline: of azaleas, associated with several parasitic nematodes, 47
---: of live oaks in Virginia, possibly caused by Endothia parasitica, 269
---: of oak and sycamore, associated with Clitocybe olearia (possible progression of wood decay organisms involved in decline), 870
---: of peach trees: possibly associated with root damage from parasitic watermolds during excessive soil moisture, Georgia, 279; survey of losses in Georgia orchards, 1965, 859
Diaporthe phaseolorum caulivora: 228
--- --- sojae: 228
Dieback: of Eucalyptus marginata, association of Phytophthora cinnamomi with, 943; survey of northern hardwoods in upper Michigan, 1954-1964, 483
Dioscorea spp.: 875
Diplodia gossypina, 138
--- tumefaciens: first record on lombardy poplar and hybrid cottonwood, 50
Diseases: in Kentucky in 1964, 450; of pea in Washington and Oregon in 1964, 235; reduction in yield of cotton caused by, 307; of snap beans in the Pacific Northwest in 1964, 637; (virus) of corn in the United States in 1964, 918
Ditylenchus dipsaci: 898; 991; onion bloat, factors affecting control by fumigants, 33
Dodder: agent of transmission for crinkle leaf disease of pokeweed, caused by unidentified virus, 1002; found on St. Johnswort and English ivy in California, 1015; transmission of Exocortis virus of citrus by, 189
Dogwood: Colletotrichum gloeosporioides infected stems and leaves in Mississippi, first report of Colletotrichum damage on dogwood, 351
Dothidella ulei: two races on several rubber tree clones, 12
Dothistroma pini: 124
Downy mildew: B strain, distribution on lima bean in 1964, 447; of hops, factors affecting spread in an arid climate, 247
Droop: Coreopsis flower, associated with unidentified toxicogenic insect, 964
Dutch elm disease: distribution in Kansas in 1964, 359; economic impact in Michigan, 830; found in Atlanta, Georgia for first time, 801; possible resistance in the European and Asiatic species of elm on University of Illinois campus, 552; transmission through root grafts reduced two-thirds with Vapam, 106
Dwarf bunt: prevalent on Gaines wheat in northern Idaho and eastern Washington in 1964, possibly due to race D-3 or an unidentified race, 401
Dwarfmistletoe: first report on foxtail pine, found in California, 647

Elm: Dutch: distribution in Kansas in 1964, 359; economic impact in Michigan, 830; found in Atlanta, Georgia for first time, 801; possible resistance in the European and Asiatic species of elm on University of Illinois campus, 552; transmission through root grafts reduced two-thirds with Vapam, 106
Endothia parasitica: 269
Endria inimica: 556
Enzyme: role of polyphenoloxidase as a factor involved in resistance of cacao to Phytophthora palmivora, 319
Epidemics: local, of fungus leaf spots on 'Gaines' wheat in Washington in 1964, 242

Ergot: test results from Nigeria indicate that fungus is more likely to infect sterile florets of grain sorghum but is not cause of host sterility, 680
Eriogonum fasciculatum: 1002
Erwinia amylovora: 769
--- dissolvens: associated with corn stunt virus disease in Georgia in 1964, 207
Erysiphe cichoracearum: 756
--- graminis f. hordei: 775
--- --- f. sp. tritici: 837
--- polygoni: 198; 787
Eucalyptus marginata: jarrah dieback, possiblity that Phytophthora cinnamomi is a contributory cause, Australia, 943
European larch canker: none found during resurvey of 2826 trees in Massachusetts, 1965, 834
Eutypa armeniacae: (imperfect stage, Cytosporina sp.), 178

Fagus grandifolia: dieback of, 483
Fertilizer: effect of 2 rates on bacterial wilt incidence in 5 solanaceous plants and peanut in newly cleared Georgia soil, 586; effect on soil microflora as indicated by fungi and bacteria isolated from roots of cotton seedlings, United Arab Republic, 1013; tolerances of Picea spp. to DBCP and a high fertilizer salt level in the soil, 72
Fig: 596
Fir (see Abies and Pseudotsuga)
Flag smut: varietal resistance in Kentucky bluegrass, 1017
Flaming: propane gas: for excellent control of mint rust, 393; for control of mint wilt, 581
Flooding: of cranberry bogs, and low dissolved oxygen concentrations failed to control nematode populations, 517
Florida: 382; 536; 742; 832; 1016; 1018
Fomes annosus: first record on Abies fraseri, 360; found in jack pine plantation in Nebraska, 456
--- pini: first record on Abies fraseri, 360
Frangipani: Cercospora plumeriae (leaf spotting fungus) found on, first report of this fungus in the United States (Florida), 729
Fungi: Ascomycetes, Basidiomycetes, and Fungi Imperfecti isolated from mycorrhizal roots of sweetgum, 869; effect of crop rotation and fertilizers on soil microflora as indicated by fungi and bacteria isolated from roots of cotton seedlings, 1013; fairy ring fungus, Marasmius oreades, causes damage to roots of turfgrasses, 571; isolated from diseased lentil seedlings in Washington, 1963-64, believed to be first report in United States of lentil diseases caused by fungi, 660; phycomycetous, association with peach tree decline in Georgia, 279; present in both normal and discolored heartwood of sugar maple in Ontario, 771; soil-inhabiting pathogenic, effect of Streptomyces rimosus and its filtrates on, in vitro, 159; western-grown pearl millet seed contains fewer destructive fungi than seed grown in Georgia, 487
Fungicides: bioassay method for detection of activity in vascular system of trees, 949; control of anthracnose and white spot of turnip greens, 419; for control of 13 fungi from peanuts, modified paper disk bioassay method used to evaluate 18 fungicides for use in the field and in storage, 928; several gave promising control of anthracnose, scab, and soil rot of cucumber, 29; effectiveness of some new fungicide formulations for control of pecan scab in Oklahoma, 285; evaluated as seed and seedling protectants and seedling eradicants of safflower rust, 623; evaluation of, for volatility, toxicity and specificity using a double Petri dish diffusion technique, 398; greenhouse evaluations of 55 compounds for control of pea root rot, 793; spray tests with new and standard fungicides for control of pecan scab, 703
---: Bordeaux mixture best of 4 tested for control of needle blight of Austrian pine, 124; 3-4-50 Bordeaux rapidly killed liverwort in tobacco plant beds in Kentucky, 590
---: Botran, excellent control for Sclerotinia wilt on green beans in Washington when sprayed on lower portion of vines, 357
---: captan and malachite green used in selective agar for isolation of Fusarium, 114; captan and zineb for control of conifer damping-off, 504
---: 2,6-dichloro-4-nitroaniline (DCNA) in the hydrocooler water gave excellent control in graded peaches of postharvest decays caused by Rhizopus stolonifer in New Jersey, 300; warm DCNA reduced decay caused by Penicillium digitatum infections of lemons during storage, 715
---: Dithane M-45 and Dyrene used with dimethyl sulfoxide (DMSO) for control of Helminthosporium diseases of Kentucky bluegrass, 810

---: Dowicide B and Nurelle, heat treatments equally effective in controlling Fusarium disease of gladiolus corms in storage, 435
---: four maneb sprays (3 containing zinc) compared for control of cucumber scab, 312
---: Merculin best of several tested for control of Alternaria disease of mango in India, 405
---: mercury compounds, excellent control of Fusarium patch disease of bentgrass turf with new experimental and commercial fungicides, 923
---: PCNB: and Botran, effectiveness in preventing white rot of onion in the field tested by an improved laboratory assay, 546; comparison of application methods for control of Sclerotinia bulborum in iris bulbs, 443; comparison of residual activities of 5 PCNB formulations for control of soil-borne pathogens, 60; effect of PCNB-treated soil on the cotton seedling disease complex under field conditions, 21; PCNB, maneb, and captan, tolerance of Rhizoctonia solani in temperature and serial transfer studies, 500; testing application methods for best control of soil-borne Sclerotium rolfsii on iris with PCNB, 149
---: Phenylmercuric acetate, residue studies on apple fruit sprayed for control of apple scab, 568
---: Phytoactin in spore germination tests, as a foliar dip or in lanolin applied to galls, did not affect fruiting by Cronartium fusiforme in vitro, 698
---: Puratized Agricultural Spray gave good control of Phomopsis blight of eastern redcedar seedlings at 3 rates used, 529
---: tetrachloroisophthalonitrile and 2, 3-5, 6-tetrachloronitroanisole compared with established turf fungicides for control of Helminthosporium and rust diseases of Kentucky bluegrass, 323
---: Vapam reduced transmission of Dutch elm disease through root grafts by two-thirds, 106
Fusarium spp.: 138; 660; isolation from soil with selective agar using captan and malachite green, 114
--- decemcellulare: 979; (imperfect stage of Calonectria rigidiuscula) macrospores induce greenpoint gall on cacao, (perfect stage may lose this property), 564
--- lateritium: 835
--- nivale: 923
--- oxysporum: 412; 928
--- --- f. sp. cepae: 522
--- --- f. gladioli: 435
--- --- f. lycopersici: 536
--- --- f. perniciosum: 971
--- --- f. sesami: 383
--- --- f. tracheiphilum: 673
--- --- f. vasinfectum: 852; 999
--- patch disease: of bentgrass turf, excellent control with new experimental and commercial fungicides, 923
Fusarium solani: 507; associated with wilting of tomato varieties resistant to F. oxysporum in Israel, 741
--- --- and F. decemcellulare: "overgrowth" of Malayan crop plants grown in infested soil correlated with gibberellin production by the fungi, 979
--- --- f. phaseoli: 630
--- wilt: of sesame, testing for resistance in South Carolina for future use, should this disease become a serious threat, 383
Fusicladium effusum: 285; 703
Fusicoccum canker: of European mountain ash, previously unreported disease found in Illinois in 1963, 251

Ganoderma sp.: (formerly referred to as G. lucidum), 1011
Gardenia: leaf spot found on plants imported from Alabama and Florida into Canada, first report of Pestalotia on gardenia, 832
Garlic: seed cloves, shortened pre-soak period at elevated temperature followed by standard hot water treatment eradicated stem and bulb nematode, 898
Georgia: 148; 207; 215; 279; 322; 487; 586; 859
Ginger: ornamental, first report of bacterial wilt of, found in Hawaii, 378
Gladiolus: heat treatments as effective as chemical dips in arresting Fusarium disease in storage when applied to corms beforehand, however further treatment immediately before planting is necessary, 435
Gloeocercospora sorghi: 91
Gloxinias: Myrothecium rot, pathogenicity affected by humidity, 78
Gnomonia caryae var. pecanae: 148
Grain smut: of jowar millet, assessment of yield loss in India, 164
Grape: 596; leaf roll virus, effect on mineral content of leaf tissue at 3 potassium levels, 36
Grasses: bentgrass, Fusarium patch disease of turf, excellent control with new experimental and commercial fungicides, 923
---: creeping bentgrass: stripe smut

damage, 710; varietal reaction to stripe smut, 268
---: Japanese lawngrass, Puccinia zoysiae on, previously reported only from Orient, found in Maryland and Missouri, 185
---: Kentucky bluegrass: use of dimethyl sulfoxide (DMSO) with 2 turf fungicides for control of Helminthosporium diseases, 810; resistance of 7 varieties to stem rust, 708; and smooth bromegrass, hosts for wheat striate mosaic virus for 7 weeks in the greenhouse, 556; susceptibility of some varieties and strains to root-knot nematodes, 89; tetrachloroisophthalonitrile and 2, 3, 5, 6-tetrachloronitroanisole compared with turf fungicides for control of Helminthosporium and rust diseases of Merion and common varieties, 323; varietal reaction to Septoria leaf spot, 802; varietal resistance to flag smut, 1017
---: lemon grass, new smut caused by Tolyposporium sp. in India, 916
---: 'Merion' bluegrass, urea effective treatment for stripe smut control, 361
---: ryegrass, annual, southern blight in Mississippi, 408
---: St. Augustine grass, parasitism and pathogenicity of Trichodorus proximus to, 259
---: turfgrasses: damage caused by cyanogenic compounds produced by Marasmius oreades, 571; melting-out of, possible fungus - nematode disease complex, 519
---: zoysia grass, rust of 3 species found in Florida for first time, 2 of which are new hosts in the United States, 382
Greenpoint gall: of cacao, induced by imperfect stage but not by perfect stage of Calonectria rigidiuscula, 564
Guar: Alternaria leafspot, identity of pathogen, symptoms, and host-specificity, 605; anthracnose of, caused by Colletotrichum dematium f. truncata, 607; hot water treatments reduced seed-borne pathogens sufficiently to be of practical value in research with infected seeds, 895
Gymnosporangium juniperi-virginianae: 578

Hardwoods: northern, 10 year survey of dieback of yellow birch, beech, red maple, and sugar maple in Upper Michigan, 483
Hawaii: 378
Heat treatments: as effective as chemical dips in arresting Fusarium disease of gladiolus in storage when applied to corms beforehand, however further treatment immediately before planting is necessary, 435; fail to inactivate barley stripe mosaic virus in seed, 696; hot water and hot air, for control of postharvest decay of red raspberries, 783; supplement refrigeration for retarding postharvest decay of strawberries, 619
Helicotylenchus dihystera: 875
Helminthosporium spp.: 810
--- diseases: of Kentucky bluegrass, use of dimethyl sulfoxide (DMSO) with 2 turf fungicides for control, 810
--- leaf spot and crown rot: usefulness of tetrachloroisophthalonitrile and tetrachloronitroanisole for control on Kentucky bluegrass, 323
--- setariae: 487
--- sorokinianum: culture filtrates of isolates, and other pathogenic genera inhibit Puccinia graminis var. tritici on 'Marquis' wheat, 371; toxic to urediospores of stem rust and not toxic to host plants, 280
--- turcicum: 516
--- vagans: 519
Herbicides: 4, 6-dinitro-o-sec-butylphenol controlled mint rust, 393; pine trees near power substations damaged by urea herbicides, 970
Heterodera cruciferae: 900; 903
--- cyperi: cyst nematode parasite of nut-grass, males necessary to complete reproductive cycle, 5
--- glycines: 747
--- schachtii; 900; 903
Hops: downy mildew, factors affecting spread in an arid climate, 247; wilt of, Verticillium dahliae and V. albo-atrum in Oregon, 538
Hoplolaimus proporicus: 875
Hosts: check list for parasitic phanerogams in Hawaii in 1963, 583
Hot water treatments: of guar seed at 54°C for 1/2 hour, effect on several seed-borne pathogens, 895; modification of standard treatment eradicated Ditylenchus dipsaci in seed garlic cloves, 898; reduced postharvest decay of unethylened oranges as much or more than standard fungicide, 463
Humidity: factor involved in Myrothecium rot of gloxinias, 78
Hydrocotyle sibthorpioides: (pennywort group, used as ground cover), greenhouse studies on nematocidal control

of root-knot nematodes, 861
Hypoxylon canker: of aspen, known southern distribution extended in Arizona, 480
--- pruinatum: 480
Hysteroneura setariae: 827

Idaho: 401
Illinois: 210; 251; 552
Immunity: to bean yellow mosaic virus in 7 cowpea cultivars, 879
Impietratura: virus disease of citrus, present status in Cyprus, 111
India: 58; 263; 405; 916; 917
Insect: toxicogenic, possible cause of Coreopsis flower droop, 964
Insecticides: Bayer 25141, 29493, and 37289 controlled root-knot nematodes on tomatoes, 603; carbaryl toxicity symptoms on soybeans, 168; dimethoate used as a nematocide for chemotherapeutic control of root-knot disease of tomato, 352
Internal mold: of pimiento, effect of fruit shape on occurrence, 157
Iodine test: for rapid detection of psorosis virus in leaves of sour orange, 18
Iowa: 339
Iraq: 792
Iris: black slime controlled by broadcast, furrow treatments or bulb dip treatments with PCNB, 443; testing application methods for best control of soil-borne Sclerotium rolfsii with PCNB, 149
Israel: 34; 741
Ivy: English, dodder found on, in California, 1015

Jarrah: (see Eucalyptus)
Juglans hindsii: etiology of blackline in, tests indicate that no virus is involved, 954
--- microcarpa: new host for Microstroma juglandis, found in New Mexico, 417
Juniperus virginiana: comparison of Phomopsis-blighted and non-blighted seedlings for field survival and growth, 121; Phomopsis blight of, good control with Puratized Agricultural Spray, 529
Jute: susceptibility of 10 varieties to 4 root-knot nematodes in Nigeria, 729

Kansas: 359
Kentucky: 450
Key: for preliminary diagnosing of virus diseases of corn in the field and laboratory, 654
Korthalsella complanata: check list of hosts found in Hawaii in 1963, 583

Larch (see Larix)
Larix spp.: (mainly L. decidua), no evidence of European larch canker found in a resurvey of 2826 trees in Massachusetts, 1965, 834
--- occidentalis: new host for Phomopsis "carrot-top" - "spike-top" disease, 102
Leaf spot: of cardamon, new disease caused by Sphaceloma sp. in India, 917; Cercospora, on peanut, correlation between explosive increase in infection and relative humidity, 975; of gardenia, found on plants imported from Alabama and Florida into Canada, first report of Pestalotia on gardenia, 832; of guar, caused by Alternaria cucumerina, symptoms and host-specificity, 605; Selenophoma and Septoria, local epidemic on 'Gaines' wheat in Washington, 242; Septoria, varietal reaction of Kentucky bluegrass to, 802; of soap-nut trees in India, new disease, 263; zonate, serious disease on sorghum-sudangrass hybrids in Alabama in 1964, 91
Leaf structure: directly related to varietal susceptibility of rice to blast disease, 170
Lentils: root and stem rots caused by 8 fungi in Washington, believed to be first report in United States of lentil diseases caused by fungi, 660
Liriodendron tulipifera: Cylindrocladium scoparium on, 797; Fusarium canker of, first report in South Carolina, 507
Liver spot: of pecan, found in Georgia, first occurrence east of Mississippi, 148
Liverwort: in tobacco beds in Kentucky, killed rapidly with 3-4-50 Bordeaux, 590
Longidorus elongatus: 509
Loose smut: failure to build up in 14 varieties of winter wheat when naturally inoculated over a period of 5 years, 239
Loquat: Septoria eriobotryae found on, in Miami, Florida, first report in the United States, 1016
Lotus spp.: Phomopsis blight, host range and testing for disease resistance, 724
Louisiana: 495

Macrosiphum avenae: 687; 692
--- rosae: 495
Malaya: 778; 979; 1011
Mango: 596; new disease caused by Alternaria tenuissima, symptomatology of

disease, pathogenicity of fungus, and efficacy of some fungicides on, 405
Maple (see Acer)
Marasmius oreades: produces cyanogenic compounds that damage roots of turfgrasses, 571
Marigold: cape, new host for Plasmopara halstedii, found in Iowa, 940; (Tagetes minuta) a cover crop that reduces root-knot nematode populations in the soil found susceptible to bacterial wilt, thus not advised for use in Pseudomonas solanacearum infested fields, 847
Maryland: 987
Massachusetts: 314; 834
Melampyrum lineare: Cronartium stalactiforme in Quebec, 909
Meloidogyne spp.: 729
--- hapla: 352
--- incognita: 822
--- incognita acrita: 723; susceptibility of some bluegrass varieties and strains to, 89
--- --- incognita: 861
Melting-out: of turfgrass, possible fungus - nematode disease complex, 519
Methods and techniques: abutilon whitefly successfully reared in the greenhouse and single viruliferous adults are capable of transmitting sweetpotato yellow dwarf virus, 429; aerial versus ground surveys for detection of oak wilt in West Virginia, 173; artist's airbrush inoculation method efficient in increasing infection in property studies of oat mosaic virus, 553; atomizer for inoculating plants with spore-oil suspension, 455; bioassay, to detect activity of fungicides translocated through vascular system of trees, 949; Colorado Soil Temperature tanks give control within ± 1°C for soil-borne pathogen study, 452; device for rapid and uniform inoculation of plants with bacteria that require wounds, 878; disposable, time and space-saving plastic growth pouches allow plants to be grown under aseptic conditions in research with soil-borne plant pathogens, 852; double Petri dish diffusion chamber used to evaluate fungicides for volatility, toxicity and specificity, 398; Freon-113 as a dispersal medium for urediospores of Puccinia graminis var. tritici, 268; inoculation and development of rust on peanuts grown in the greenhouse, 191; iodine test for rapid detection of psorosis virus in leaves of sour orange, 18; laboratory evaluation of seed treatments to control rice seed rot, 736; modified dew duration recorder used in monitoring moisture of cotton plant parts susceptible to disease, 193; quantitative recovery of nematodes with a modified Baermann funnel influenced by temperature, 662; selective agar for isolation of Fusarium with use of captan and malachite green, 114; serological detection of halo blight, specific and rapid, 297; standardization of artificial inoculation of cucumber with Colletotrichum lagenarium for obtaining uniform results, 649; tetrazolium (TTC) used in method for estimating degree of resistance or susceptibility of various citrus rootstocks to Phytophthora root rot, 423; two new tools for field, and nursery and greenhouse nematode subsampling, 265
Mexico: 417
Michigan: 483; 527; 830; 1016
Microstroma juglandis: first report on little walnut, found in New Mexico, 417
Millet: jowar, use of two mathematical formulae to correlate incidence of grain smut with loss in yield, 164
Mimosa: Fusarium wilt found along Mississippi coast, 971
Minerals: content of K, Ca, and Mg in leaf roll virus-infected grape leaf tissue, 36
Minnesota: 527; 646
Mint: rust, excellent control by applying propane flame or alkanolamine salts of 4,6-dinitro-o-sec-butylphenol to soil surface at shoot emergence, 393
---: wilt, propane flaming kills Verticillium dahliae in peppermint stubble, 581
Mississippi: 84; 753; 971
Missouri: 213
Mulberry: mosaic disease of, in India, 981
Mushroom: fly agaric, temporary toxicity of extracts to the chrysanthemum foliar nematode resembles effect of narcotics on the nervous system, 1017
Mutants: of normally sensitive strains of Pencillium digitatum, Type A which cannot sporulate in absence of biphenyl, and Type B which can, 965
Mycorrhizae: Ascomycetes, Basidiomycetes, and Fungi Imperfecti isolated from, in roots of sweetgum, 869
Myzus persicae: 1018

Nebraska: 269; 456
Nematocides: reaction of Brussels sprouts to soil fumigation for control of sugar-

beet and cabbage root nematodes, 903
---: DBCP: gave effective control of root-knot nematode on Hydrocotyle sibthorpioides in the greenhouse, 861; gave excellent control of plant-parasitic nematodes in citrus when used in irrigation water in Sudan, 598; most effective of several in transplanting water of Brussels sprouts for control of sugarbeet and cabbage root nematodes, 900; one application controlled nematodes associated with blue spruce for 2 years, 69; successfully controlled root-lesion nematode disease of roses in containers and plants showed significant growth responses, 982; tolerance of Picea spp. to DBCP in heavily fertilized soil, 72
---: Zinophos: attempts to improve growth of Radopholus similis-infected citrus with under-tree drenches, 865; bare-root dips in Zinophos and Bayer 25141 give good control of root-knot, lesion, and burrowing nematodes in nursery plants, 51
Nematodes: apparently not correlated to 4 viruses in Florida citrus groves, 610; 6 of 8 chemical amendments used in soil culture of alfalfa did not harm Xiphinema americanum population levels, 994; effect of Tylenchorhynchus brevidens alone or with Olpidium brassicae (fungus) on yield of hard red winter wheat, 806; non-transmission of certain strawberry viruses by, 509; possible fungus - nematode complex associated with melting-out of turfgrass, 519; quantitative recovery from soil samples with modified Baermann funnel influenced by temperature, 662; soil fumigation improved seedling emergence, vigor, and yield of sugarbeet regardless of degree of control, 600; two new tools for field, and nursery and greenhouse subsampling, 265
---: burrowing, attempts to improve growth of infected citrus with under-tree drenches of Zinophos, 865
---: cabbage root: on Brussels sprouts, effectiveness of nematocides in transplant water, 900; effectiveness of soil fumigation, 903
---: chrysanthemum foliar, temporary toxicity of extracts of Amanita muscaria resembles effect of narcotics on nervous system, 1017
---: citrus-root, found in citrus groves in Iraq for first time, 1965, 792
---: cotton root-knot, potato losses in South Carolina due to, 723
---: cyst, of soybean, reaction of sugarbeet varieties to, 747
---: new species, 6 previously unreported species associated with Florida citrus 610
---: plant parasitic: associated with serious decline of azaleas, 47; controlled in blue spruce nursery planting with DBCP, 69; in established citrus groves, DBCP applied in irrigation water gave excellent control, 598; on raspberries in the eastern United States, survey results suggest possible significance in production, 987; 11 species found on yam over entire survey area in Nigeria, 875; survey in established orchards of newly-reclaimed sandy areas of United Arab Republic, 596
---: root-knot, lesion, and burrowing controlled in nursery plants by chemical bare-root dips, 51; root-lesion and root knot, tobacco varieties used to show differential response to, 822
---: root-knot: on Hydrocotyle sibthorpioides (used as ground cover), greenhouse studies on nematocidal control, 861; susceptibility of jute strains to 4 Meloidogyne spp. in Nigeria, 729; on tomato, controlled with dimethoate, 352, with organophosphates, 603
---: root-lesion: reaction of white clover and 5 other crops to, 856; on roses, controlled in plants growing in containers with drench treatments of emulsifiable DBCP, 982; on roses, more severe in light sandy soils at higher temperatures (85°F) than in heavier soil at lower temperatures (65°F), 849
---: stem and bulb: first record on Penstemon digitalis, description of injury to, 991; in seed garlic cloves, shortened pre-soak period at elevated temperature followed by standard hot water treatment for eradication, 898
---: stylet-bearing, 5 species not controlled by flooding cranberry bogs for 2 years, 517
---: sugarbeet, on Brussels sprouts: effectiveness of nematocides in transplant water, 900; effectiveness of soil fumigation, 903
Nephotettix impicticeps: (=N. bipunctatus), 127; 778
New diseases: blister of sweetpotato, cause unknown, 97; crinkle leaf of soybean produced by unreported bacterial pathogen, 396; tomato condensed top virus, new graft transmissible virus in Sudan, 599;

unidentified virus causes spur type growth of Bing cherry, may be valuable in controlling tree size or may present complications, 910
---: in India: blight and leaf spot are two new Phyllosticta diseases of two types of soap-nut trees, 263; leaf spot of cardamon caused by undescribed species of Sphaceloma, 917; leaf spot rot on several unrelated host plants, 58; smut of lemon grass caused by Tolyposporium sp., 916
---: in the United States: corn showing stunting, leaf mottling, and abnormal tassels and ears is associated with sap-transmissible virus, Illinois, 210; Fusicoccum canker of European mountain ash, previously unreported disease found in Illinois in 1963, 251; Phytophthora root and crown rot of vetch in Mississippi, 84
New distribution: barley yellow dwarf virus on oats in Israel, 34
---: Cercospora plumeriae found in the United States (Florida) for the first time, 729
---: Dutch elm disease found in Atlanta, Georgia for first time, 801
---: first known case of McIntosh apple trees naturally infected with russet ring virus in the United States, 314; first report of lentil diseases caused by fungi in the United States, found in Washington, 1963-64, 660
---: Fusarium canker, of tulip poplar in South Carolina, 507
---: --- wilt, of cotton in Arizona, 179
---: liver spot disease of pecan in Cordele, Georgia, 148
---: oak wilt in Texas and Arkansas: 1007
---: Phytophthora heveae found in United States for first time in forest soil in North Carolina and Tennessee, 233
---: Puccinia zoysiae: on Japanese lawngrass, new to North America, 185; on Zoysia japonica in Florida, 382; on Zoysia tenuifolia and Z. matrella in the United States, found in Florida, 382
---: Septoria eriobotryae found in United States for first time, (on loquat in Miami, Florida), 1016
---: willow blight in the central United States (Michigan), 1016
New hosts: Armillaria mellea on several, found in California, 812
---: Articularia quercina var. minor on Quercus dunnii, found in New Mexico, 417
---: Colletotrichum gloeosporioides on flowering dogwood in Mississippi, 351
---: Diplodia tumefaciens on lombardy poplar and hybrid cottonwood, 50
---: Ditylenchus dipsaci on Penstemon digitalis, this is the first record although disease was observed in North America (Virginia) 5 years ago, 991
---: dwarfmistletoe on foxtail pine, found in California, 647
---: Fomes annosus, Polyporus schweinitzii and Fomes pini on Abies fraseri, North Carolina, 360
---: halo blight on pea, found in Pacific Northwest, 854
---: Microstroma juglandis on Juglans microcarpa, found in New Mexico, 417
---: Pestalotia langloisii on gardenia, found on plants imported from Alabama and Florida into Canada, 832
---: Phomopsis pseudotsugae on western larch, 102
---: Phytophthora wilt of Tephrosia vogelii, found in Puerto Rico, 676
---: Plasmopara halstedii on cape marigold, 940
---: Pseudomonas solanacearum on Tagetes minuta (a nematode resistant marigold used as cover crop), 847
---: Pseudomonas solanacearum, on Hedychium flavum, H. coronarium, and H. gardnerianum, found in Hawaii, 378
---: Scleroderris canker on subalpine fir, found in Colorado, 882
---: Sphaerotheca fuliginea on Zinnia elegans, new host in India, 756
---: tobacco ringspot virus in birdsfoot trefoil, first record of natural occurrence, 855
New Mexico: 417; 904
Nicotiana glutinosa (see also Tobacco): testing effect of some growth-retarding compounds on tobacco mosaic virus, 775
Nigeria: 680; 729; 875
Nitrogen: study with 12- and 16-hour light periods for effect on susceptibility of sesame to bacterial leaf spot, 119
North Carolina: 233; 360; 797
Nursery stock: chemical bare-root dips for control of nematodes, 51
Nut-grass: cyst nematode parasite of, males necessary to complete reproductive cycle, 5
Nutrients: carbaryl toxicity symptoms similar to those of nutrition problems on soybean, 168; effect of leaf roll virus on K, Ca, and Mg content of grape leaf tissue at 3 levels of K and influence on Mn uptake, 36

Oak: first report of Articularia quercina var. minor on Gambel oak, found in New Mexico, 417; unsuccessful study to determine progress of Poria and Polyporus decay in 10 years, 81
---: decline: Clitocybe olearia associated with, (possible progression of wood decay organisms involved in decline), 870; possibly caused by Endothia parasitica, in Virginia, 269
---: fusiform rust, resistance increases in leaves as they mature, 972
---: wilt: aerial surveys in West Virginia, 173; distribution extended in Oklahoma (first report in Craig County), 954; distribution in Tennessee in 1964, 68; reported distribution in 616 counties in 20 States in 1964, 1007; spread in Nebraska, 269
Oats: barley yellow dwarf virus: found in Israel for first time, 34; variations in nature of interactions among virus aphid vectors, and hosts, and a variation of epidemiology in the field, 692
---: crown rust, variability greater in physiologic races of aecial isolations than of uredial isolations from collections on buckthorn, 132
---: mosaic virus, soil-borne, artist's airbrush inoculation method efficient in increasing infection for property studies, 553
---: southern blight, in Mississippi, 408
---: stem rust, resistance of Ukraine and derivatives, (possibly related to crown rust resistance of Ukraine), 403
Ohio: 615
Oil: spray, number 3 paraffin oil mixed with spreader-sticker Triton B-1956 inhibited field spread of potato virus A, 557
Oklahoma: 285; 806; 954
Olpidium brassicae: 806
Onion: bloat, factors affecting control by soil fumigants, 33
---: Fusarium oxysporum f. sp. cepae, field trials of 14 varieties for reaction to, western New York, 522
---: Stemphylium botryosum conidia violently released when conidial apparatus subjected to rapid vapor decrease, 1006
---: white rot: assay to determine protective potential of soil chemicals against specific host-pathogen-environment combination, 546; testing for an artificial inoculation method for use in breeding for resistance, 760
Ontario: 771
Oregon: 284; 538; 636; 684
"Overgrowth": symptoms in Malayan crop plants as a result of Fusarium infection, 979

Pacific Northwest: 275; 637; 837; 854
Palm: basal stem rot, results of root inoculations of oil palm seedlings in pots with Ganoderma sp., 1011
Panagrolaimus sp.: 519
Papaya: papaya mosaic and ringspot viruses isolated from diseased plants in Florida, 742
Pea: diseases in Washington and Oregon, 1964, 235; halo blight (Race 2a), found in Pacific Northwest for first report on this host, 854; legume viruses, severity measured by plant growth reduction, 627; root rot, greenhouse evaluations of 55 soil fungicides for control, 793; studies on translocation and titer increase of three virus isolates, 24; USDA plant introduction lines of Pisum sativum tested for reactions to alfalfa mosaic, clover yellow mosaic, and pea streak viruses and to powdery mildew, 787
---: cow: bean yellow mosaic virus, 7 cultivars immune to, 879; Choanephora pod rot, disease reaction of varieties and breeding lines, 347
Peach (see also Prunus): bacterial canker in the southeastern States, 701; comparison of effect of adding 2,6-dichloro-4-nitroaniline to dump tank, wet brusher and hydrocooler for control of Rhizopus stolonifer rot of graded harvested peaches, 300; decline, survey to determine extent of losses due to, Georgia, 1965, 859; tree decline in Georgia, possibly associated with root damage from parasitic watermolds during excessive soil moisture, 279
Peanut: laboratory evaluation of 18 fungicides against 13 fungi found in peanuts for use in the field and in storage, 928; survey for extent of pod and kernel invasion by Aspergillus flavus at digging time, Alabama, Georgia, North Carolina, 931
---: bacterial wilt, low disease incidence in plants growing in infested soil, 586
---: leafspot, correlation between explosive increase in infection and relative humidity, 975
---: rust, inoculation and development on 7 varieties grown in the greenhouse, 191
Pear: diseases in Kentucky in 1964, 450

---: bark measles, occurrence in Colorado Bartlett orchards in 1962, 392
---: fire blight, testing comparative resistance of Stewart with other Bartlett clones under controlled conditions, 769
Pearl millet: western-grown seed contains fewer destructive fungi than seed grown in Georgia, 487
Pecan: freeze damage causes abnormal flowering, 146
---: liver spot disease found in Georgia, first occurrence east of Mississippi, 148
---: scab: effectiveness of some new fungicide formulations for control, 285; spray tests with new and standard fungicides for control, 703
Pellicularia filamentosa (Rhizoctonia solani): 711
Penicillium spp.: 928
--- digitatum: 332; 965; warm DCNA reduced decay of lemons during storage, 715
Penstemon digitalis: first record of Ditylenchus dipsaci on, description of injury to, 991
Pepper: bacterial spot, several chemicals tested in culture and in the field for control, 748
---: collar-rot; black Puerto Rican species tested for susceptibility to pathogen to determine suitability as rootstocks, 550
---: fruit rot, 48 varieties and breeding selections of sweet and hot peppers tested for field susceptibility to Alternaria tenuis, 886
Peronospora tabacina: 154; 367
Persimmon: wilt, possible use as a tree eradicant for weed persimmons, 789
Pestalotia langloisii: 832
Petunia: suspected transmission of exocortis virus of citrus to, by dodder, 189
Phomopsis juniperovora: 529
--- loti: 724
--- pseudotsugae: (=Phomopsis strobi), 102
Photoperiod: varying effect on susceptibility of 2 sesame varieties to 2 bacterial leaf spots, in relation to the addition of supplemental nitrogen, 119
Phyllosticta sp.: 263
Phytophthora spp.: 490; collar-rot of pepper, 550
--- cinnamomi: association with jarrah dieback in Australia, 943
--- citrophthora: 423
--- cryptogea: 753
--- drechsleri: 293
--- erythroseptica: believed to be first report of Phytophthora on vetch in United States, 84
--- heveae: found in United States for first time in forest soil in North Carolina and Tennessee, 233
--- megasperma: 753
--- palmivora: 135; 319; 565
--- parasitica: 327; 423
--- --- var. nicotianae: 215; wilt of Tephrosia vogelii, first report on this host, 676
--- phaseoli: (strain "B"), 447
Picea spp.: tolerances to DBCP and a high fertilizer salt level, 72
--- pungens: nematodes in nursery controlled with DBCP, 69
Pimientos: effect of fruit shape on occurrence of internal mold, 157
Pinus spp.: eastern and western gall rusts on Jack pine, distribution in the Lake States, 527; progeny tests indicate little difference in susceptibility between lines from rust-infected and noninfected loblolly pine parent trees, 872
--- balfouriana: first report of dwarf-mistletoe on, found in California, 647
--- banksiana: Cronartium stalactiforme in Quebec, 909; Fomes annosus killing trees in Nebraska, 456
--- elliottii: fusiform rust, cultures of gall tissue were maintained in an apparently viable condition for 5 months, 616
--- --- var. elliottii: damage by urea herbicides near power substation, 970; and P. taeda, occurrence of Polyporus tomentosus sporophores on basal stem cankers caused by Cronartium fusiforme in Georgia plantations, 322
--- nigra: needle blight of, Bordeaux mixture best of 4 fungicides tested for control, 124
--- radiata: damping-off of, promising treatments for control in South Australia, 504
--- strobus: Cylindrocladium scoparium on, 797; negative control of blister rust in North Carolina and Tennessee with cycloheximide, 342
Piricularia oryzae: 170
Plasmopara halstedii: on Dimorphotheca sinuata, 940
Pleurotus ostreatus and P. ulmarius: 986
Plum (see also Prunus): bacterial canker in the southeastern States, 701; rusty blotch, virus-like disorder of Japanese-type plums, incidence increasing

in California, 109
Podosphaera leucotricha: 876
Pokeweed: crinkle leaf, caused by undescribed virus transmitted by dodder from desert shrubs in California, 1002
Pollacia saliciperda: (imperfect stage of Venturia saliciperda), 1016
Polyporus fissilis and P. hispidus: study of progress of decay in artificially inoculated oaks, 81
--- schweinitzii: first record on Abies fraseri, 360
--- tomentosus: 322
Poplar (see Populus)
Poplar, yellow-, (see Liriodendron)
Populus spp.: additional hosts of Diplodia tumefaciens (=Macrophoma tumefaciens), 50
--- deltoides: blackstem of, studies indicate that it is an infectious disease and that susceptibility may be a heritable trait, 867
--- tremuloides: Hypoxylon canker of, known southern distribution extended in Arizona, 480
Poria ambigua: study of progress of decay in artificially inoculated oaks, 81
Postharvest decays: of graded peaches, rot caused by Rhizopus stolonifer controlled by addition of 2,6-dichloro-4-nitroaniline to hydrocooler water, 300
Potato: alfalfa mosaic virus, isolate from potato induced enation symptoms in tobacco, 684
---: cotton root-knot nematode, heavy losses in South Carolina due to, 723
---: decay of seed pieces, effect of storage temperature, variety, storage time, and chemical treatment, 638
---: virus A, field spread inhibited by oil, 557
Potentilla spp.: 2 species severely infected by Xanthomonas fragariae in the greenhouse, 491
Powdery mildew: of apple, 3 compounds screened at Tree Fruit Experiment Station have mildewcidal properties, 876; of barley, effect of some growth-retarding compounds on the fungus, 775; of cabbage, inheritance of resistance controlled by single dominant gene, 198; of pea, USDA plant introduction lines of Pisum sativum tested for reaction to, 787; of Rubus henryi, 814; of wheat, importance on semidwarf varieties such as Gaines in the Pacific Northwest, 837; of Zinnia elegans caused by Sphaerotheca fuliginea rather than Erysiphe cichoracearum, first report from India, 756
Pratylenchus brachyurus: 822
--- scribneri: 856
--- vulnus: 849; 982
Pruning: marked effect in reducing Phytophthora pod rot of cacao, 257
Prunus spp. (see also common names): release of virus-indexed budwood from Interregional repository, 187; shock symptoms of necrotic ringspot and prune dwarf, 288
--- salicina: plum rusty blotch steadily increasing in California, 109
Pseudomonas sp.: 380
--- phaseolicola: 297; 591; 854
--- sesami: 119
--- solanacearum: 378; 847
--- syringae: 591; 701
Pseudoperonospora humuli: 247
Pseudotsuga menziesii: recurrence of Cytospora canker in Colorado, 481
Puccinia arachidis: 191
--- carthami: 440; 623
--- coronata avenae: 132
--- erianthi: 764
--- graminis: 708; 775
--- --- agrostis: 323
--- --- f. sp. avenae: 403
--- --- var. tritici: 63; 268; 280; 37'
--- helianthi: 433
--- hordei: 575
--- menthae: 393
--- recondita: 222
--- --- tritici: 308
--- striiformis: 275; 335; 636
--- zoysiae: 185
Puerto Rico: 550; 676
Pythium spp.: 736
--- ultimum: (or a similar type) cause of pre-emergence damping-off of soybean in Minnesota, 646

Quarantine: significance of host-pathogen relationship between Araucaria excelsa and Cryptospora longispora is discussed, 656
Quebec: 909

Races: of Dothidella ulei on several rubber tree clones, Race 1 from Guatemala and Race 2 from Costa Rica, 12; Race 3 of Pseudomonas solanacearum indigenous on newly cleared land in Georgia, 586
---: bunt: D-3 or unidentified race of Tilletia controversa on 'Gaines' wheat in the Pacific Northwest, 401; screening of spring wheat varieties for resistance to common bunt races, 839; screening of winter wheat varieties for resistance to common bunt races, 843
---: Fusarium: Race 1 of cowpea-wilt

causes chrysanthemum wilt, not aster-wilt Fusarium as previously reported, 673; Fusarium oxysporum f. lycopersici biotype (possibly Delray) pathogenic to Race 1-resistant tomato stocks found in 2 locations in Manatee County, Florida, 536
---: rust: of barley leaf rust, identification of 4 races including new Race 53, 575; North American 1965 set of supplemental differential wheat varieties for leaf rust race identification (analytical key), 308; racial variability greater in aecial than in uredial collections of oat crown rust on buckthorn, 132; of Puccinia graminis in the United States in 1963, 63
Radopholus similis: 865
Ranunculus asiaticus: biological properties of a virus isolated from, compared with ranunculus mosaic and anemone mosaic previously described from California and England, 449
Raspberry: chlorotic leaf spot (bushy dwarf) evidence of transmission in pollen and seed, 230; postharvest decay, testing for control with hot water dips, hot air treatments, chemical dips, and fumigation with SO_2, 783; survey results suggest that plant nematodes may be significant in production in the eastern United States, 987
Relative humidity: compilation of literature sources for maintaining constant RH with 4 types of solutions, 948; effect on infection with peanut leafspot, 975
Repository: budwood, virus-indexed Prunus materials released for research and industry as of August 1964, 188
Reproduction: males necessary for completion of reproductive cycle in cyst nematode parasite of nut-grass, 5
Residues: of 5 PCNB dust and granular formulations in the soil, 60
Rhizoctonia sp.: 660
--- solani: 29; 60; 235; chemical seed treatments tested for control of cotton seedling damping-off, 720; fungicide tolerance influenced by temperature and serial transfer in fungicide-treated soil, 500; (Pellicularia filamentosa) differences in reaction of 5 cotton varieties to 3 isolates suggest existence of physiologic specialization, 711; tolerant isolates possible reason for cotton seedling damping-off in PCNB-treated soil, 21; twenty-one isolates tested for virulence and factors affecting disease severity on alfalfa and red clover, 512
Rhizopus sp.: 619; 783; 928
--- stolonifer: 300
Rhode Island: 750
Rhododendron: crown gall, two records in North America, 360
Rhopalosiphum maidis: 202; 665; 687
--- padi: 34; 687; 692
Rhubarb: abscission of petiole bases in stored stalks, 959
Rice: blast, structure of leaves directly related to varietal susceptibility, 170
---: "penyakit merah" (red disease), in Malaysia believed to be caused by a virus, and possibly related to tungro virus of rice in the Philippines, 778
---: seed rot: technique for laboratory evaluation of 7 seed treatments for control, 736; laboratory tests of seed treatments for control, 1964, 738
---: "tungro" disease (="accep na pula") is caused by a virus transmitted by Nephotettix impicticeps, 127; relationship to other rice virus diseases, studied with antiserum produced in rabbits, 305
Root grafting: potential source of error in apple indexing, 39
Rootstock: relation of selection and of tree age in scion overgrowth disorder, 7
Rosa sp.: veinbanding virus, Macrosiphum rosae as a vector, first report in Louisiana, 495
Rose: root-lesion nematode disease, controlled in plants growing in containers with drench treatments of emulsifiable DBCP, 982; more severe in light sandy soil at higher temperatures (85°F) than in heavier soil at lower temperatures (65°F), 849
Rot: in thinned sweetgum sprout clumps apparently not hazardous to remaining stems for at least a decade, 986
---: basal, Fusarium, reaction of 14 onion varieties in field trials, 522
---: black, of crucifers, evaluation of phenacridane chloride as a disinfectant and seed protectant, 439
---: boll, of cotton, observations in South Carolina, 138
---: buckeye, of tomato, influence of maturity of fruit, source of isolate, temperature, and incubation period on infection and disease development, 327
---: cob: of open-pollinated corn, positive correlation between low pH of cob tissues and resistance to infection, 246; Nigrospora cob rot, of corn in

the United Arab Republic, reaction of 4 varieties to, 244
---: collar-: of black pepper, Puerto Rican species tested for susceptibility to pathogen to determine suitability as rootstocks, 550; hypocotyl, of bean seedlings, possible connection with bacterium, 134
---: crown: of iris, testing application methods for best control of soil-borne Sclerotium rolfsii with PCNB, 149; Phytophthora, of alfalfa seedlings in Mississippi, 753
---: fruit, of peppers, 48 varieties and breeding selections of sweet and hot peppers tested for field susceptibility to Alternaria tenuis, 886
---: leaf spot, new sclerotial disease on several unrelated hosts in India, 58
---: Myrothecium, of gloxinias, pathogenicity affected by humidity, 78
---: pod: of cacao, behavior of pathogen in soil, 135; Choanephora, of cowpea, disease reaction of varieties and breeding lines to, 347; Phytophthora, of cacao: effect of spacing and pruning on, 257; role of polyphenoloxidase as a factor involved in resistance, 319
---: root, of pea, greenhouse evaluations of 55 soil fungicides for control, 793
---: ---, Fusarium, of pea, 235; testing Phaseolus species for resistance, 630
---: ---, Phytophthora: of alfalfa seedlings and mature plants, in Mississippi, 753; of citrus, tetrazolium (TTC) used in test for resistance or susceptibility, 423; of safflower and sugarbeet, pathogenicity and growth rates of P. drechsleri isolates, 293; of soybean in Minnesota, 2 new resistant varieties, 490; of vetch in Mississippi, 84
---: --- and butt, on planted slash and loblolly pine in Georgia, 322
---: --- and stem: comparison of cultural characteristics of several Cylindrocladium scoparium isolates and their pathogenicity to yellow-poplar, white pine, and Fraser fir in North Carolina, 797; of lentils, believed to be first report in United States of lentil diseases caused by fungi, 660; of sesame and castorbean, caused by Thielaviopsis basicola, 256
---: seed, of rice: laboratory evaluation of seed treatments to control, 736; laboratory tests of seed treatments in 1964 for control, 738
---: soil, destructive disease of cucumber in Florida, promising control, 29
---: southern stem, effect of some growth-retarding compounds tested on bean seedlings, 775
---: storage, of gladiolus corms, controlled as effectively by heat treatments as by chemical dips, 435
---: white, of onion: assay to determine protective potential of soil chemicals against specific host-pathogen-environment combination, 546; testing for an artificial inoculation method for use in breeding for resistance, 760
Rubber trees: reaction of several clones to two races of Dothidella ulei, 12
Rubus henryi: (used in virus indexing of raspberry plants) found to be susceptible to powdery mildew in the greenhouse, 814
Russet: stem cavity, of apples, 225
Rust: Cronartium stalactiforme on jack pine in Quebec, 909; diseases of Kentucky bluegrass, usefulness of tetrachloroisophthalonitrile and tetrachloronitroanisole for control, 323; of mint, excellent control by applying propane flame or alkanolamine salts of 4,6-dinitro-o-sec-butylphenol to soil surface at shoot emergence, 393; on peanut, inoculation and development in the greenhouse, 191; 18 safflower introductions may represent several genetic sources of resistance to both foliage and seedling phases, 440; in slash and loblolly pine, association of sporophores of Polyporus tomentosus with rust stem canker in Georgia plantations, 322; of sugarcane, in Tanganyika, 764; of sunflowers, in the Mississippi Delta, 433; of wheat, Freon-113 as a suspending medium for dispersal of urediospores, 268; on Zoysia japonica, new to North America, 185; on Zoysia tenuifolia and Z. matrella new to the United States, and of Z. japonica new to Florida, 382
---: blister, of white pine, not controlled in the Southeast by cycloheximide, contrary to results in the Pacific Northwest, 342
---: cedar apple, 12 apple clonal rootstock varieties studied for susceptibility to, 578
---: crown, of oats, racial variability greater in aecial than in uredial collections on buckthorn, 132
---: eastern and western gall, of jack pine, distribution in the Lake States, 527
---: fusiform: cultures of gall tissue from slash pine were maintained in an apparently viable condition for 5

months, 616; of loblolly pine, tests indicate little difference in susceptibility between progeny lines from infected and noninfected parent trees, 872; of oak, as leaves mature they become increasingly resistant to infection by either uredia or telia, 972; Phytoactin in spore germination tests, as foliar dip, or in lanolin applied to galls, did not affect fruiting by Cronartium fusiforme in vitro, 698
---: leaf: of barley, survey for new pathogenic strains for breeding improvement research, 575; North American 1965 set of supplemental differential wheat varieties for race identification (analytical key), 308; of wheat, identification of resistance combinations by differential temperature effects, 222
---: safflower seedling, efficacy of antifungal substances for control, 623
---: stem: inhibited by culture filtrates of isolates of Helminthosporium sorokinianum and other pathogenic genera on 'Marquis' wheat, 371; Helminthosporium sorokinianum toxic to urediospores of, 280; resistance of 7 Kentucky bluegrass varieties to, 708; 'Ukraine' oat resistance to, (possibly linked to crown rust resistance of Ukraine), 403; of wheat: effect of some plant growth-retarding compounds on, 775; physiologic races in the United States in 1963, 63
---: stripe-: evaluation of resistance to head infection in 5 Pacific Northwest wheats, 335; oversummering at high elevations in the Pacific Northwest, 275; of wheat and barley, possibly first report of stem infection, 636
Rye: winter, possible reservoir for overwintering wheat striate mosaic virus in North Dakota, 556

Safflower: isolation of alfalfa mosaic and lettuce mosaic viruses from necrosis affected plants, 541
---: root rot, pathogenicity and growth rates of Phytophthora drechsleri isolates from safflower and sugarbeet, 293
---: rust, 18 introductions may represent several genetic sources of resistance to both foliage and seedling phases, 440
---: seedling rust, evaluation of antifungal substances for control, 623
Scab: evaluation of the cucurbita for resistance, list of screened accessions supplied upon request, 388
---: of cucumber: destructive disease in Florida, promising control, 29; fungicide sprays for control, 312
---: of pecan: effectiveness of some new fungicide formulations for control, 285; spray tests with new and standard fungicides for control, 703
Schizaphis graminum: 687
Scleroderris canker: of subalpine fir in Colorado, previously unreported on this host, 882
Sclerotinia bulborum: 443
--- camelliae: 258
--- sclerotiorum: 660
--- white mold: of pea, 235
Sclerotium bataticola: 928
--- cepivorum: 546; 760
--- rolfsii: 58; 149; 408; 775
Scolytus multistriatus: 801
Scutellonema spp.: 875
Seed: abortion in citrus fruits affected with stubborn disease, 668
--- borne, unidentified bacterium isolated from watermelon plant introduction seedlings, 818
---: transmission, virus disease (Johnson grass strain of sugarcane mosaic) is seed-borne in both field and sweet corn, 468
--- treatments: effect on cotton seed from different harvest dates and storage periods, 720; evaluation of fungicides for control of seedling safflower rust, 623; evaluation of phenacridane chloride as a disinfectant and seed protectant for control of black rot of crucifers, 439; H_2SO_4 prescribed by quarantine regulations detrimental to germination of Norfolk Island pine seed, 656; of Pinus radiata seed with captan and zineb for control of damping-off, 504; technique for laboratory evaluation of fungicides to control rice seed rot, 736; tested in the laboratory for control of seed rot of rice, 738; unsuccessful attempt to free barley seed of stripe mosaic virus by heat treatments, 696
Seedling disease complex: of cotton, 3 Alternaria spp. should be considered important in, New Mexico, 904
Selenophoma donacis: 242
Septoria eriobotryae: first report in the United States, (found on loquat in Miami, Florida), 1016
--- macropoda var. septulata: 802
--- tritici: 242
Serum: anti-Pseudomonas phaseolicola, for detection of halo blight, 297; anti-, to tungro virus of rice obtained by intravenous and intramuscular

injections in rabbits, 305
Sesame: bacterial leaf spot, susceptibility of 2 varieties to pathogens affected by light-period, nitrogen treatments, 119
---: Fusarium wilt; testing for resistance in South Carolina for future use, should this disease become a serious threat, 383
---: root and stem rot, susceptibility to Thielaviopsis basicola, 256
Smut: of lemon grass, new disease caused by Tolyposporium sp. in India, 917
Soap-nut tree: blight and leaf spot are two new Phyllosticta diseases of two Sapindus spp. in India, 263
Soil: selective agar for isolation of Fusarium with use of captan and malachite green, 114; behavior of Phytophthora palmivora in, 135; Phytophthora heveae found in forest soil of North Carolina and Tennessee for first report in United States, 233; effect of type on root-lesion nematode disease of roses, 849; effects of Streptomyces rimosus and its filtrates on several pathogenic soil fungi, 159; temperature tank improvement, 452
---: treatments: factors affecting control of onion bloat by soil fumigants, 33; fumigation for control of cabbage root and sugarbeet nematodes, response of Brussels sprouts to, 903; for reduction of sulfur deficiency in broccoli in the field, 891; sugarbeet seedling emergence, vigor, and yield better when grown in fumigated soil, 600
Solutions: compilation of inorganic salts, sulfuric acid, bases, and carbohydrates for maintaining constant relative humidity, 948
Sophora japonica: trunk and branch canker of young trees caused by Fusarium lateritium, 835
Sorghum: ergot of, suggested control by pollination in Nigeria, 680
---: Verticillium wilt, successful use of disposable, time and space-saving plastic growth pouches which allow plants to be grown under aseptic conditions in studies of penetration by V. albo-atrum into roots, 852
---: zonate leafspot, serious on sorghum-sudangrass hybrids in Alabama in 1964, 91
South American leaf disease: of Para rubber trees, 12
South Carolina: 258; 507; 723
Soybean: carbaryl toxicity symptoms, 168
---: crinkle leaf, new disease produced by unreported bacterial pathogen, 396
---: Diaporthe phaseolorum, effect of variants on germination and growth in New Jersey, 228
---: Phytophthora root rot, 2 new resistant varieties -- Harosoy 63 and Chippewa 64, 490
---: Pythium pre-emergence damping-off, in Minnesota, 646
Spacing: planting distances affect incidence of Phytophthora pod rot of cacao, 257
Sphacelia sorghi: 680
Sphaceloma sp.: 917
Sphacelotheca sorghi: 164
Sphaerotheca fuliginea: 756
--- humuli: on Rubus henryi, 814
Spike-top: stage of Phomopsis disease on western larch, 102
Spinach: beet mosaic virus in leaves usually confined to parenchyma and only rarely enters the phloem, 375
Spray: relation of application schedules to mercury residue of apple fruit, 568
Spruce (see Picea)
Squash (see Cucurbits)
Stem canker: of guar seedlings, significantly reduced by hot water treatment of seed at 54°C for 1/2 hour, 895; of soybean, effect on germination and growth in New Jersey, 228
Stemphylium botryosum: violent conidia release occurs with a rapid decrease in vapor pressure, 1006
St. Johnswort: dodder found on, in California, 1015
Storage: decay of potato seed pieces, effect of temperature, time, variety, and chemical treatment, 638; petiole base abscission in stored rhubarb stalks, 959
Strawberries: certain chemical and heat treatments supplement refrigeration for retarding postharvest decay, 619; diseases in Kentucky in 1964, 450; five viruses were not transmitted by Oregon populations of three nematodes, 509; hybrid selections, rapid spread of mottle and mild yellow-edge viruses, in Oregon, 284; mottle virus, Acyrthosiphon porosum and Aphis gossypii as vectors in Louisiana, 495
Streptomyces rimosus: and its filtrates, effect on several pathogenic soil fungi, 159
Stripe smut: of creeping bentgrass, reaction of 9 varieties and 11 experimental varieties to, 268; of 'Merion' bluegrass, urea found to be an effective treatment for control, 361; damage on 'Pennlu' creeping bentgrass, 710
Subsampling: two new tools for field and

nursery and greenhouse nematode subsampling, 265
Sudan: 599
Sugarbeet: nematodes: reaction to the soybean cyst nematode, 747; seedling emergence, vigor, and yield better when grown in fumigated soil, whether nematode free or infested, 600
---: seedling blight and root rot, pathogenicity and growth rates of Phytophthora drechsleri isolates from safflower and sugarbeet, 293
---: viruses: beet mosaic virus, effect of variety of different strains (which differ in virulence) on root yields, 961; infection by a strain of cucumber mosaic virus restricted to leaf parenchyma, 375
Sugarcane: rust, in Tanganyika, 764
---: viruses: infection with mechanically transmissible corn virus, maize dwarf mosaic (possibly a strain of sugarcane mosaic), 757; sugarcane mosaic virus: recovered from naturally infected corn in California, 204; study of transmission by the rusty plum aphid, 827
Sulfur deficiency: of field-grown broccoli (first report), description of symptoms in field and greenhouse, 891
Sulfur dioxide: 3 year study shows SO_2 in the air causes most damage to late spring plants, 911
Sunflower: rust, in the Mississippi Delta, 433
Sweetgum: Ascomycetes, Basidiomycetes, and Fungi Imperfecti isolated from mycorrhizal roots of, 869; decay in thinned sprout clumps apparently not hazardous to remaining stems for at least 10 years, 986
Sweetpotato: blister, a new disease, cause unknown, 97
---: yellow dwarf virus: abutilon whitefly successfully reared in the greenhouse and single viruliferous adults are capable of transmitting the virus, 429; susceptibility of several varieties to, 425
Sycamore: Clitocybe olearia associated with decline of, 870

Tanganyika (East Africa): 764
Temperature: effect on decay of potato seed pieces during storage, 638; effect on fungicide tolerance in Rhizoctonia solani in vitro, 500; effect on root-lesion nematode disease of roses, 849; identification of wheat leaf rust resistance combinations by differential temperature effects, 222; influence on initiation of crown gall in woody hosts, 734; influence on quantitative recovery of nematodes from soil samples important in field evaluation of nematocides, 662; improved soil tanks for soil-borne pathogen study in Colorado, 452
Tennessee: 68; 233
Texas: 999; 1007
Tephrosia vogelii: Phytophthora wilt (first report), differences in susceptibility among plant introductions suggest the existence of possible genetic resistance, 676
Thielaviopsis basicola: 256
Tilletia caries: 42; 839; 843
--- controversa: 401
--- foetida: 839; 843
Tobacco (see also Nicotiana): diseases in Kentucky in 1964, 450
---: alfalfa mosaic virus, enation symptoms induced by isolates from snap beans and potatoes, 684
---: bacterial wilt, Race 3 of Pseudomonas solanacearum slightly virulent to, 586
---: black shank, in flue-cured, spread and control in Georgia since 1959, 215
---: blue mold: in Israel, 154; comparison of European and Beltsville isolates of Peronospora tabacina, 367
---: liverwort in plant beds in Kentucky killed with 3-4-50 Bordeaux, 590
---: nematodes, several varieties used to show differential response to root-lesion and root-knot, 822
---: ringspot virus, effect of mechanical inoculation of various plant parts on symptom expression on several cucurbits and beans, 54
Tolyposporium sp.: 916
Tomato: diseases in Kentucky in 1964, 450
---: bacterial canker: description of cotyledonary symptoms, 634; and spotted wilt in the west-central Great Plains, 386
---: bacterial wilt: Race 3 of Pseudomonas solanacearum highly virulent to, 586
---: buckeye rot, influence of maturity of fruit, source of isolate, temperature, and incubation period on infection and disease development, 327
---: condensed top virus, new graft transmissible virus in Sudan, 599
---: Fusarium wilt: of varieties resistant to F. oxysporum possibly caused by F. solani, 741; probable spread

in Florida of biotype that is pathogenic to Race 1-resistant varieties, 536
---: potato-Y virus, precise information on the effect of this aphid-borne virus on yield of tomatoes in south Florida, 1018
---: root-knot nematodes: controlled with 3 organophosphate insecticides, 603; greenhouse tests with dimethoate for control, 352
Transmission: field spread of apple chlorotic leaf spot virus between adjacent trees suggests direct mechanical leaf to leaf or branch transmission, 958; of chlorotic leaf spot (bushy dwarf) of raspberry in pollen and seed, evidence of, 230; of exocortis virus of citrus by dodder, 189; of sugarcane mosaic virus by the rusty plum aphid, 827
Trialeurodes abutilonea: 425; 429
Trichocladium canadense: 771
Trichodorus allius: 509
--- christiei: 47
--- proximus: parasitism and pathogenicity to St. Augustine grass, 259
Tylenchorhynchus brevidens: 806
--- claytoni: 47
Tylenchulus semipenetrans: 792

United Arab Republic: 596; 1013
Urocystis agropyri: 1017
Ustilago striiformis: 268; 361; 710
--- tritici: 239

Vectors associated with: barley yellow dwarf virus, of oats: -- Macrosiphum avenae and Rhopalosiphum padi, 692; -- Rhopalosiphum padi, 34
---: corn virus -- corn leaf aphid, 202
---: maize dwarf mosaic virus, of corn -- corn leaf aphid, 665
---: mosaic disease, of mulberry -- Aphis gossypii, 981
---: mottle virus: of strawberry -- Acyrthosiphon porosum, and Aphis gossypii, 495; and mild yellow-edge virus of hybrid strawberries -- strawberry aphids, 284
---: "penyakit merah" (red disease), of rice -- Nephotettix impicticeps, 778
---: potato-Y virus, on tomatoes -- green peach aphid, 1018
---: sugarcane mosaic virus: -- rusty plum aphid, 827; in corn -- corn leaf, pea, green peach, and cotton aphids, 204
---: "Tungro" disease, of rice -- rice green leafhopper, 127
---: veinbanding virus, of Rosa sp. -- Macrosiphum rosae, first report from Louisiana, 495
---: wheat striate mosaic, of winter rye -- leafhoppers, 556
---: yellow dwarf virus: of barley -- oat bird-cherry aphid, English grain aphid, corn leaf aphid, and greenbug, 687; of sweetpotato -- abutilon whitefly: 425; 429
---: yellow mosaic disease, on mung bean -- sweetpotato whitefly, 166
Venturia inaequalis: 568
--- saliciperda: (imperfect stage - Pollacia saliciperda), 1016
Verticillium albo-atrum: 538; 660; 852; 999
--- dahliae: 538; 581
Vetch (Vicia angustifolia): Phytophthora root and crown rot in Mississippi, believed to be first report of Phytophthora disease on vetch in United States, 84
Virginia: 269; 665
Virus, Virus disease: biological properties of a virus from Ranunculus asiaticus compared with ranunculus mosaic and anemone mosaic previously described from California and England, 449; dwarfness in Bing cherry caused by, may be valuable means of controlling tree size or may present complications, 910; of leguminous forage plants in Rhode Island, 750; pitting of citrus varieties in Israel, 936; release of virus-indexed Prunus budwood from interregional repository, 187; testing susceptibility of crambe to, 74
---: alfalfa mosaic: caused no apparent damage to safflower plants also infected with necrosis causing lettuce mosaic, 541; isolates from snap beans and potatoes induced enation symptoms in tobacco, Oregon, 684; USDA plant introduction lines of Pisum sativum tested for reaction to, 787
---: of apple: chlorotic leaf spot, field spread between adjacent trees suggests direct mechanical leaf to leaf or branch transmission, 958; fruit russet ring, first known case of naturally infected McIntosh apple trees in the United States, 314; latent viruses indicated on 'Hopa' crab grafts in Oregon, 15; second season symptoms of latent virus infection possibly result from root grafting, 39; leaf flecking, of McIntosh apple in Massachusetts, 314
---: of barley: stripe mosaic, unsuccessful attempt to free seed of virus by heat treatments, 696
---: ---: yellow dwarf: isolated from naturally infected oats, first report in Israel, 34; of oats, variations in

nature of interactions among virus, aphid vectors, and hosts, and a variation of epidemiology in the field, 692; survey for predominating strains in New York for 1964, 687
---: of bean: curly top (Ruga verrucosous), differential tolerance of 52 snap bean varieties, 475; yellow mosaic: 7 cowpea cultivars immune to, 879; of moong (mung bean), effect of vector control in India, 166
---: beet mosaic, in sugarbeets, effect of variety of different strains (which differ in virulence) on root yields, 961
---: of citrus: exocortis, transmission to citron by dodder, and suspected transmission to petunia, 189; impietratura, present status in Cyprus, 111; psorosis infection rapidly detected by iodine test of leaves, 18; stubborn disease, excessive seed abortion in affected fruit, 668; tristeza, xyloporosis, psorosis, and exocortis, apparently not correlated to nematodes in Florida citrus groves, 610
---: clover yellow mosaic, USDA plant introduction lines of Pisum sativum tested for reaction to, 787
---: of corn: in Arkansas, 202; in Iowa, mechanically transmitted, 339; in the United States in 1964, 918; sap-transmissible virus associated with new mottle-stunt disease of corn in southern Illinois, 210; tentative key for preliminary field and laboratory decisions, 654
---: ---: maize dwarf mosaic: distribution and yield loss in Ohio in 1964, 615; possibly a strain of sugarcane mosaic, isolated from corn, tested for effect on sugarcane, 757; possibly a strain of sugarcane mosaic virus, occurrence and transmission in Virginia, 665; variation in resistance among 93 sweet corn varieties suggests the presence of resistance genes in some varieties, 478
---: ---: stunt disease: status in Georgia in 1964, 207; symptoms and distribution in Missouri in 1964, 213
---: ---: sugarcane mosaic: recovered from naturally infected corn, milo maize, and Johnson grass in California, 204; (Johnson grass strain), seed-borne in both field and sweet corn, 468
---: cucumber mosaic, juice-inoculated beet leaves restricted to parenchyma tissue and spreads slowly, 375
---: Cucumis 2C, in bottlegourd, effect of certain chemotherapeutic agents on, 493
---: grape leaf roll, effect on mineral content of leaf tissue at 3 potassium levels, 36
---: lettuce mosaic, isolated from safflower, associated with severe necrosis, 541
---: of mulberry, mosaic disease, in India, 981
---: oat mosaic, soil-borne (Marmor terrestre), artist's airbrush inoculation method efficient in increasing infection for property studies, 553
---: papaya mosaic (MM) and ringspot (DR) isolated from diseased papaya plants in Florida, 742
---: of pea: BYMV, PEMV, AMV, PSV, and CYMV, severity in peas measured by reduction of plant growth, 627; pea streak I and II, and pea mosaic II, translocation and titer studies, 24; pea streak, USDA plant introduction lines of Pisum sativum tested for reaction to, 787
---: pokeweed crinkle leaf, caused by undescribed virus transmitted by dodder from desert shrubs in California, 1002
---: potato A, field spread inhibited by oil, 557; potato-Y, precise information on effect of this aphid-borne virus on yield of tomatoes in south Florida, 1018
---: of Prunus: necrotic ringspot and prune dwarf symptoms in several Prunus spp., 288
---: ---: cherry: necrotic ringspot, latent in Washington orchards, 143; necrotic rusty mottle, a disease with similar symptoms associated with bacterial infection, 380; "short stem", 253; sour cherry yellows, latent in Washington sweet cherry orchards, 143
---: ---: plum rusty blotch, virus-like disorder of Japanese-type plums in California, 109
---: of raspberry, chlorotic leaf spot (bushy dwarf), evidence that disease is pollen-seedborne, 230
---: of rice: "penyakit merah" (red disease), in Malaysia, believed to be caused by a virus, and possibly related to tungro virus of rice in the Philippines, 778; tungro disease, antiserum, produced by immunization of rabbits, used to test relation of tungro to other virus diseases of rice in the Philippines, 305; tungro disease caused by a virus transmitted

by the rice green leafhopper, 127
---: squash mosaic disease, study of in Egypt, 218
---: of strawberry: Macrosiphum rosae, Acyrthosiphon porosum, and Aphis gossypii as vectors in Louisiana, 495; mottle and mild yellow-edge, rapidly spreading in Oregon, 284; non-transmission by 3 nematodes in Oregon, 509
---: sugarbeet mosaic, in spinach leaves largely confined to the parenchyma and does not readily enter the phloem to infect healthy growing shoots, 375
---: sugarcane mosaic: transmission by the rusty plum aphid, 827
---: of sweetpotato: yellow dwarf (feathery mottle), adaptation of abutilon whitefly for laboratory use, single viruliferous adults can transmit virus, 429; yellow dwarf (Nanus ipomoeae), susceptibility of several varieties, 425
---: of tobacco: mosaic, effect of some growth-retarding compounds on, tested on Nicotiana glutinosa, 775
---: ---: tobacco ringspot: effect of mechanical inoculation of various plant parts on symptom expression on several cucurbits and beans, 54; first report of natural occurrence in birdsfoot trefoil, 855
---: of tomato: condensed top, new graft transmissible virus in Sudan, 599; spotted wilt virus, in west-central Great Plains, previously unreported from Wyoming, 387
---: watermelon mosaic 1 and 2, separated and identified in cucurbit specimens from production areas in 6 States and Mexico, 532
---: of wheat: streak-mosaic, new source of resistance, 813; striate mosaic, overwintering under North Dakota conditions, 556

Walnut (see Juglans)
Washington: 242; 357; 401; 660; 765; 891
Watermelon (see Cucurbits)
Weather: factors influencing SO_2 phytotoxicity in New Jersey, 911; freeze damage causes abnormal flowering of pecan, 146; modified dew duration recorder used in monitoring moisture on cotton plant parts susceptible to disease, 193
Wheat: bunts: effective control with HCB, PCNB, and TCNA in the Pacific Northwest, 42; dwarf bunt prevalent on Gaines in northern Idaho and eastern Washington in 1964, possibly due to Race D-3 or an unidentified race, 401; resistance screening of 244 spring varieties from the Uniform Regional Spring Wheat Nursery to 4 key races of bunt from 1954-62, 839; resistance screening of 102 winter varieties from the cooperative uniform wheat varietal trials to 7 key races of bunt since 1956, 843
---: leaf rust: identification of resistance combinations by differential temperature effects, 222; North American 1965 set of supplemental differential wheat varieties for leaf rust race identification, 308
---: loose smut, failure of naturally disseminated inoculum to build up in 14 varieties, 239
---: powdery mildew, assay on semi-dwarf varieties in 1964 in Pacific Northwest, economic importance, pathogenic races, and host resistance, 837
---: stem rust: effect of some plant growth-retarding compounds on, 775; Helminthosporium sorokinianum toxic to urediospores of, 280; inhibition by culture filtrates of isolates of Helminthosporium sorokinianum and other pathogenic genera, 371; physiologic races in the United States in 1963, 63
---: rust, Freon-113 as a suspending medium for dispersal of urediospores of Puccinia graminis var. tritici, 268
---: Selenophoma and Septoria leaf spots, local epidemic on 'Gaines' and Burt-7 varieties in Washington, 1964, 242; streak-mosaic virus, new source of resistance, 813
---: stripe-rust: evaluation of resistance to head infection in 5 Pacific Northwest varieties, 335; possibly first report of stem infection, Oregon, 636; oversummering at high elevations in the Pacific Northwest, 275
---: yield depressing effect on hard red winter wheat varieties resulting from land infested with the nematode Tylenchorhynchus brevidens, either alone or in association with the fungus Olpidium brassicae, 806
White spot: (pale spot), of turnip greens, fungicidal control, 419
Willow: blight, found in Michigan for first report from central United States, 1016
Wilt: bacterial: of Hedychium spp., caused by Pseudomonas solanacearum found in Hawaii, first report, 378; identification and persistence of an indigenous race of Pseudomonas

solanacearum in a soil in Georgia, 586; on South American marigold (a cover crop that reduces root-knot nematode populations in the soil), first report on this host, 847
---: Fusarium: of chrysanthemum, caused by Race 1 of cowpea-wilt Fusarium, not by aster-wilt Fusarium as previously reported, 673; of cotton, first occurrence in Arizona, 179; of mimosa, found along Mississippi coast, 971; pathogenicity studies of 3 forms of F. oxysporum which cause similar symptoms in several varieties of alfalfa, 412; recurrence in Florida of F. oxysporum f. lycopersici biotype pathogenic to Race 1-resistant tomato stocks, 536; of tomato varieties resistant to F. oxysporum possibly caused by F. solani, 741; Fusarium and Verticillium wilts in South Plains area of Texas correlated with soil type, 999
---: oak: aerial surveys for detection of, in West Virginia, 173; distribution extended in Oklahoma (first report in Craig County), 954; distribution in Tennessee, 1964, 68; reported distribution in 616 counties in 20 States in 1964, 1007; spread in Nebraska, 269
---: persimmon, possible use as a tree eradicant for weed persimmons, 789
---: Phytophthora, of Tephrosia vogelii, first report on this host, 676
---: Sclerotinia, on green beans in Washington, excellent control with Botran when sprayed on lower portion of vines, 357
---: Verticillium: of hops, in Oregon, 538; of mint, controlled by propane flaming of stubble after harvest, 581
Wisconsin: 516; 527; 591
Witches' broom: first report on little walnut and on Gambel oak, found in New Mexico, 417
Wyoming: 386; 813

Xanthomonas sp.: 380
--- campestris: 439
--- cyamopsidis: 895
--- fragariae: two Potentilla spp. severely infected in the greenhouse, 491
--- sesami: 119
--- vesicatoria: 748
Xiphinema americanum: 994
--- bakeri: 509

Yam: eleven species of plant-parasitic nematodes found on, in entire survey area in Nigeria, 875
Yellow-poplar (see Liriodendron)

Zinnia elegans: powdery mildew usually caused by Erysiphe cichoracearum, first report in India caused by Sphaerotheca fuliginea, 756

CROPS RESEARCH DIVISION

Prepared by Sheila D. Thompson

3/3840